

12

**EUROPEAN PATENT APPLICATION**

21 Application number: 86107199.1

22 Date of filing: 27.05.86

51 Int. Cl.<sup>4</sup>: **C 07 D 491/22**  
**C 07 D 495/22, C 07 D 513/22**  
**C 07 D 498/22, C 07 D 471/20**  
**C 07 D 471/22, A 61 K 31/435**  
**A 61 K 31/505, A 61 K 31/50**  
**A 61 K 31/535, A 61 K 31/55**  
**/(C07D491/22, 307:00, 235:00,**  
**221:00, 221:00), (C07D491/22,**  
**307:00, 307:00, 221:00, 221:00),**  
**(C07D491/22, 307:00, 231:00,**  
**221:00, 221:00)**

30 Priority: 03.06.85 US 740609

43 Date of publication of application:  
10.12.86 Bulletin 86/50

84 Designated Contracting States:  
AT BE CH DE FR GB IT LI LU NL SE

71 Applicant: MERCK & CO. INC.  
126, East Lincoln Avenue P.O. Box 2000  
Rahway New Jersey 07065(US)

72 Inventor: Baldwin, John J.  
621 Gypsy Hill Circle  
Gwynedd Valley, PA 19437(US)

72 Inventor: Huff, Joel R.  
738 Bergey Mill Road  
Ledersach, PA 19450(US)

72 Inventor: Vacca, Joseph P.  
766 Eisenhower Drive  
Telford, PA 18969(US)

72 Inventor: Young, Steven D.  
J-10 Forge Gate Apts. Snyder Road  
Lansdale, PA 19446(US)

72 Inventor: De Solms, Jane  
735 Port Indian Road  
Norristown, PA 19403(US)

72 Inventor: Guare, James P., Jr.  
R.D. 2 Kumry Road  
Quakertown, PA 18951(US)

74 Representative: Blum, Rudolf Emil Ernst et al,  
c/o E. Blum & Co Patentanwälte Vorderberg 11  
CH-8044 Zürich(CH)

54 Substituted hexahydro arylquinolizines, processes for their preparation and pharmaceutical compositions containing them.

57 Substituted hexahydro arylquinolizines and pharmaceutically acceptable salts thereof are selective  $\alpha_2$ -adrenergic receptor antagonists and thereby useful as antidepressants, antihypertensives, ocular antihypertensives, antidiabetics, platelet aggregation inhibitors, antiobesity agents, and modifiers of gastrointestinal motility.

1

TITLE OF THE INVENTION

Substituted hexahydro arylquinolizines,  
processes for their preparation and pharmaceutical  
5 compositions containing them.

BACKGROUND OF THE INVENTION

This invention is concerned with novel  
10 substituted hexahydro arylquinolizines or  
pharmaceutically acceptable salts thereof which are  
selective  $\alpha_2$ -adrenoceptor antagonists and are of  
value in conditions where selective antagonism of the  
 $\alpha_2$ -adrenoceptor is desirable for example as  
15 antidepressant, antihypertensive, ocular anti-  
hypertensive, antidiabetic, antiobesity agents,  
platelet aggregation inhibitors, and modifiers of  
gastrointestinal motility. It also relates to  
processes for preparing the novel compounds,  
20 pharmaceutical compositions comprising the novel  
compounds and to a method of antagonizing  
 $\alpha_2$ -adrenoceptors.

25

30

2291S/0976A

- 2 -

172261A

The concept that the complex clinical state of depression is linked to a functional deficiency of monoamines in the central nervous system is now widely accepted. Numerous biochemical and clinical observations support the proposal that many forms of depressive illness are associated with reductions in adrenergic activity at functionally important sites in the brain. Thus, classical antidepressive drugs, such as amitriptyline and imipramine, are believed to act by blocking the neuronal reuptake of norepinephrine and/or serotonin, thereby enhancing the availability of the monoamines as neurotransmitters.

In addition to  $\alpha_1$ -adrenergic receptors which mediate postsynaptic responses to the neurotransmitter, norepinephrine, other adrenergic receptors are present at or near sympathetic terminals. These latter receptors,  $\alpha_2$ -adrenergic receptors, form part of a negative feedback system which modulates noradrenergic neurotransmission by controlling the impulse-induced release of norepinephrine from presynaptic terminals. Activation of  $\alpha_2$ -adrenergic receptors results in a decrease in the amount of norepinephrine normally released from the nerve terminals by nerve impulses while antagonism of  $\alpha_2$ -adrenergic receptors increases norepinephrine release. Therefore, molecules that block  $\alpha_2$ -adrenergic receptors afford an alternate approach to enhancement of noradrenergic function and the treatment of depression associated with an absolute or relative deficiency of adrenergic function.

2291S/1163A

- 3 -


172261A

$\alpha_2$ -Adrenergic receptor antagonism is also associated with antidiabetic, antihypertensive, ocular antihypertensive, antiobesity, platelet aggregation inhibition activity, and modification of gastrointestinal motility.

Compounds structurally related to the novel compounds of this invention are disclosed in British Patents 1,435,573 and 2,106,909 of John Wyeth and Brother, Ltd.


#### DETAILED DESCRIPTION OF THE INVENTION

This invention is concerned with a compound of structural formula I:




I


or a pharmaceutical acceptable salt thereof, wherein Ar represents an aromatic heterocycle such as:




pyridino-




benzofuro-




thiazolo-




benzo-




imidazo-




pyrazolo-



benzothieno-



thieno-



furo-

2291S/0976A

- 4 -

17226IA

$R^1$  and  $R^2$  are independently,

- 1) hydrogen,
- 2) halo, such as chloro, bromo, or fluoro,
- 3) hydroxy,
- 5 4)  $C_{1-3}$ alkoxy, or
- 5)  $C_{1-6}$ alkyl, either straight or branched chain;

B represents a spiroheterocycle of 4-7 members with up to 2 heteroatoms one of which is bonded to the spiro carbon and if Ar is benzo that heteroatom is nitrogen wherein the members are independently  $-CH_2-$ ,  $-\underset{\substack{| \\ R^3}}{CH}-$ ,  $-\underset{\substack{|| \\ O}}{C}-$ ,  $-\underset{\substack{|| \\ S}}{C}-$ ,  $-\underset{\substack{|| \\ NCN}}{C}-$ ,  $-O-$ ,

- $-NR_3$  or  $-SO_2-$  such as, (1)spiro-4-imidazolidin-2-one, (2)spiro-4-imidazolidin-2-thione, (3)spiro-3-(1,2,5-thiadiazolidin-1,1-dioxide), (4)spiro-5-pyrazolidin-3-one, (5)spiro-5-pyrrolidin-2-one, (6)spiro-5-tetrahydrofuran-2-one, (7)spiro-5-oxazolidin-2-one, (8)spiro-4-oxazolidin-2-one, (9)spiro-3-isoxazolidin-5-one, (10)spiro-4-imidazolidin-2,5-dione, (11)spiro-4-azetidin-2-one, (12)spiro-4-(5,6-dihydro-1H-pyrimidin-2(3H)-one), (13)spiro-4-(1,3-diazin-2,6-dione), (14)spiro-4-(3,4,5,6-tetrahydro-1,3-oxazin-2-one), (15)spiro-5-(2,4,5,6-tetrahydro-1,4-oxazin-3-one), (16)spiro-5-piperazin-2,3-dione, (17)spiro-5-piperazin-3-one, or (18)spiro-5-(1,4-diazepin-7-one).


$R^3$  is 1) hydrogen,

- 2)  $-\underset{\substack{|| \\ O}}{C}-R$ , wherein R is hydrogen or  $C_{1-3}$  alkyl,
- 3)  $C_{1-6}$  alkyl, either unsubstituted or substituted with one or more of;

2291S/1163A

- 5 -

17226IA

- 5
- a) hydroxy,  
 b) carboxy,  
 c)  $C_{1-3}$ alkoxycarbonyl,  
 d) halo such as fluoro, chloro or  
 bromo,
- 10
- e)  $C_{1-3}$ alkoxy,  
 f)  $-CONR^6R^7$  wherein  $R^6$  and  $R^7$   
 are the same or different and are  
 hydrogen or  $C_{1-5}$ alkyl or joined  
 together either directly to form a  
 5-7 membered ring such as  
 pyrrolidino, or piperidino, or  
 through a heteroatom selected from  
 N, O, and S, form a 6-membered  
 heterocycle with the nitrogen to  
 which they are attached such as  
 morpholino, piperazino,  $N-C_{1-3}$   
 alkylpiperazino,
- 15
- g)  $-NR^6R^7$ ,
- 20
- h) 
- i)  $-SO_2NR^6R^7$ ; or (  
 j)  $-SO_2(C_{1-3}alkyl)$ ;
- 25
- $R^5$  is
- 1) hydrogen,  
 2)  $C_{1-6}$  alkyl, either unsubstituted or  
 substituted with one or more of  
 a)  $-OR^8$ , wherein  $R^8$  is  
 i) H, or  
 ii)  $C_{1-6}$ alkyl,  
 b)  $N(R^8)COR^8$ , or  
 c)  $CO_2R^8$ ,  
 3)  $-CO_2R^8$ , or  
 4)  $-CONR^6R^7$ .
- 30



2291S/1163A

- 7 -


17226IA

The pharmaceutically acceptable salts coming within the purview of this invention include the pharmaceutically acceptable acid addition salts. Acids useful for preparing these acid addition salts include, inter alia, inorganic acids, such as the hydrohalic acids (e.g., hydrochloric and hydrobromic acid), sulfuric acid, nitric acid, and phosphoric acid, and organic acids such as maleic, fumaric, tartaric, citric, acetic, benzoic, 2-acetoxybenzoic, salicylic, succinic, theophylline, 8-chloro-theophylline, p-aminobenzoic, p-acetamidobenzoic, methanesulfonic, or ethanedisulfonic acid.

In a preferred embodiment of this invention, Ar is  $R^1, R^2$ -benzo[b]furo-,  $R^1, R^2$ -benzo[b]-thieno-, thieno-, furo- or benzo- and B is a spiro-4-imidazolidin-2-one or spiro-4-(5,6-dihydro-1H-pyrimidin-2(3H)-one). It is further preferred that  $R^1$  and  $R^2$  be hydrogen or halo and  $R^3$  be  $C_{1-6}$  alkyl, especially methyl. It is also preferred that  $R^5$  be hydrogen or  $C_{1-6}$  alkyl.

It is most preferred that  $R^1$  and  $R^2$  be hydrogen,  $R^3$  be methyl, and  $R^5$  be hydrogen.

The novel compounds of this invention are described herein as having the configuration in which the heteroatom in ring B attached to carbon 2 and the hydrogen at 12b are trans-,




30



and it is the more preferred isomer for  $\alpha_2$ -adrenoceptor blockade activity. However, the isomers in which the heteroatom in ring B attached to carbon 2 and the hydrogen at 12b are cis are also active  $\alpha_2$ -adrenoceptor blockers and are considered to be within the scope of this invention. Each of these configurational isomers are racemates capable of being resolved into dextrorotatory and levorotatory enantiomers. This invention includes these pure enantiomers as well as all mixtures thereof, especially the racemates.

Another embodiment of this invention are the novel processes used to prepared the novel compounds.

For those compounds wherein the spiroheterocycle, B, includes a carbonyl or thiocarbonyl flanked on both sides by a heteroatom independently selected from oxygen and nitrogen, i.e. wherein B is an imidazolidin-2-one, or thione, oxazolidin-2-one or thione, or dihydro-pyrimidin-2-one or thione, or 2,6-dione a preferred process is to treat the 2,2-disubstituted quinolizine with carbonyl-diimidazole or thiocarbonyldiimidazole as represented by the following:



2291S/1163A


- 9 -

172261A

The reaction is conducted in an inert organic solvent such as benzene, toluene, chloroform, methylene dichloride, or the like or mixtures thereof at about 10 to 50°C. The temperature is not critical and the reaction is most readily conducted at room temperature. Reaction times of about 1 to about 10 hours are sufficient to complete the reaction, but longer times are not deleterious.

As an alternative to the carbonyl diimidazole and thiocarbonyldiimidazole there may be employed phosgene or thiophosgene respectively.

Similarly for those novel compounds wherein the spiroheterocycle includes a sulfonyl flanked on both sides by an oxygen or nitrogen such as a thiadiazolidin-1,1-dioxide, a 2,2-disubstituted quinolizine is treated with sulfuryl chloride as represented by the following:




This reaction is best conducted in an inert atmosphere and solvent such as chloroform, methylene dichloride, benzene, toluene or the like at about -10 to +10°C, and preferably about 0°C for about 1 to 8 hours followed by spontaneous warming to room temperature or about 20°C.

Another ring forming reaction useful for preparing the novel compounds of this invention comprises treating a quinolizin-2-one with a mixture of an organometallic such as n-butyl lithium and allyl-bis-(dimethylamino)phosphonate in an ethereal solvent such as THF, diethyl ether or 1,2-dimethoxyethane at about -50 to -20°C followed by spontaneous warming to room temperature over about 2 to 5 hours. The reaction is depicted as follows:

10

15




20

Spiro-imidazolidin-diones are prepared from 2-amino-2-alkoxycarbonylquinolizines by treatment with alkyl isocyanates depicted as follows:

25

30



2291S/1163A


- 11 -

172261A

The reaction is conducted in an inert organic solvent such as acetonitrile, dioxane, chloroform, or the like preferably at about room temperature, although temperature is not critical, for about 12 to 24 hours.

5 Spiro-pyrazolidin-ones are prepared by treating a 2-alkoxycarbonylmethylenylquinolizine with hydrazine in an inert organic solvent such as benzene, toluene or the like at about 65-100°C for about 0.5 to 3 hours. The reaction is as shown below:

10



20 Similarly, using methylhydrazine in refluxing THF for about 18 to 48 hours there is produced the corresponding 2-methylpyrazolidin-3-one.

The same ester starting material with ethylenediamine in a lower alkanol such as methanol at about room temperature for about 20-60 hours provides a spiro-5-(1,4-diazepin-7-one).

25 The same ester starting material is useful in the synthesis of spiro-isoxazolidin-5-ones by treatment with N-alkyl hydroxylamine and an alkali metal carbonate such as potassium or sodium carbonate in an inert organic solvent such as THF, diethyl ether, or the like at about reflux temperature (30-50°C) for about 12 to 48 hours.

30


2291S/1163A

- 12 -

172261A

Treatment of a 2-amino-2-aminomethyl-quinolizine with diethyloxalate results in the formation of a spiro-piperazine-2,3-dione. The reaction is conducted in the same manner as that described for synthesis of the imidazolidinones.


Treatment of a 2-chloroacetoxymethyl-2-chloroacetyl-amino quinolizine with a mixture of finely divided potassium hydroxide and neutral alumina suspended in an inert organic solvent such as benzene or toluene at about ambient temperatures for about 1 to 4 hours provides a spiro 1,4-oxazin-3-one, as shown below:



In a similar reaction a 2-trifluoroacetyl-aminomethyl-2-chloroacetyl-aminoquinolizine is treated with potassium hydroxide pellets in a lower alkanol such as methanol or ethanol at ambient temperature for about 1 to 4 hours to yield a spiro-piperazin-3-one.

Another spirocycle of this invention is the 4-membered azetidinone and it is readily prepared by treating a spiro-isoxazolidin-5-one with hydrogen in the presence of a noble metal catalyst at a slightly elevated pressure of about 20 to 80 psig. at or near

room temperature in a lower alkanol solvent for about 12 to 24 hours, or until the requisite amount of hydrogen is absorbed. The resulting amino acid is cyclized by treatment with dicyclohexylcarbodiimide (DCC). The ring size reduction is depicted as follows:



Any of the spiro-heterocycles with a secondary amino group as a member of the ring may be alkylated, or benzylated, or acylated by standard techniques well known to those skilled in the art.

In the novel method of this invention of selectively antagonizing  $\alpha_2$ -adrenergic receptors in a patient, a novel compound or pharmaceutically acceptable salt thereof, is administered in an amount ranging from about 0.01 to about 20 mg per kg of body weight per day, preferably from about 0.1 to about 10 mg per kg of body weight per day in a single dose or in 2 to 4 divided doses.

The novel compounds of this invention can be administered as the sole active ingredient or in combination with other antidepressants such as amitriptyline, imipramine or other norepinephrine or serotonin reuptake inhibitor or a monoamine oxidase inhibitor.

2291S/1163A

- 14 -

17226IA

These doses are useful for treating depression, diabetes, hypertension, ocular hypertension, abnormal platelet aggregation, obesity and abnormal gastrointestinal motility.

5           The compounds, or pharmaceutically acceptable salts thereof, of the present invention, in the described dosages, are administered orally, intraperitoneally, subcutaneously, intramuscularly, transdermally or intravenously. They are preferably  
10 administered orally, for example in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gum, or the like prepared by art recognized procedures. The amount of active  
15 compound in such therapeutically useful compositions or preparations is such that a suitable dosage will be obtained.

#### EXAMPLE 1

(2RS,12bSR)-3'-methyl-spiro(1,3,4,6,7,12b-hexahydro-  
20 benzo[b]furo[2,3-a]quinolizine)-2,4'-imidazolidin-2'-one  
Step A: Preparation of 3-Cyanomethylbenzo[b]furan

To a suspension of 2.64 gms (0.11 mole) of oil free sodium hydride in 200 ml of tetrahydrofuran (THF) was added dropwise a solution of 19.47 gms  
25 (0.11 mole) of diethylcyanomethylphosphonate in 75 mL of THF. After the H<sub>2</sub> evolution had ceased, a solution of 13.4 g (0.1 mole) of 3-(2H)-benzo[b]-furanone in 100 mL of THF was added. The solution was heated at 70°C for 1.5 hours, cooled, and poured  
30 into 500 mL of 5% HCl, and washed with ether. The ether phase was washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated to give 15.4 g of a dark oil. The product was distilled at 96-100°C/0.075 mm Hg to give 10.85 g of a yellow oil which crystallized upon standing.

2291S/1163A

- 15 -

172261A

Step B: Preparation of 2-(3-benzo[b]furanyl)ethyl-  
amine

A solution of 3.97 g (0.025 mole) of  
3-cyanomethylbenzo[b]furan in 200 mL of diethyl ether  
5 was slowly added to a refluxing suspension of 3.84 g  
(0.1 mole) of lithium aluminum hydride in 400 mL of  
ether. The reaction was heated 3 hours., cooled and  
water was slowly added. The suspension was filtered  
through a pad of filter aid and the filtrate was  
10 evaporated to give 3.2 g of oily product. The  
hydrochloride salt has m.p. 183-185°C.

Step C: Preparation of 3-(2-Formamidoethyl)benzo-  
[b]furan

15 A solution of 2.35 g (0.015 mole) of 2-(3-  
benzo[b]furanyl)ethylamine and 5 mL of ethyl formate  
was heated at 60°C for 3 hours, poured into 2N HCl  
and washed with methylene chloride which in turn was  
washed with 5% sodium hydroxide (w/v), dried  
20 (MgSO<sub>4</sub>), filtered and concentrated to give 2.70 g  
of product.

Step D: Preparation of 3,4-dihydrobenzo[b]furo-  
[2,3-c]pyridine

25 2.28 Grams (0.012 mole) of 3-(2-formamido-  
ethyl)benzo[b]furan was added to 28 g of  
polyphosphoric acid which was preheated to 100°C.  
After 1-1.5 hours, the reaction mixture was poured  
onto ice and the residues were washed with water.  
30 The polyphosphoric acid was dissolved in water,  
filtered through a pad of celite and made basic with  
concentrated ammonia. A precipitate was collected  
and dried to give 1.45 g of product, m.p. 170-171°C.



2291S/0976A

- 16 -

172261A

Step E: Preparation of (12b-SR)-1,3,4,6,7,12b-Hexahydrobenzo[b]furo[2,3-a]quinolizin-2-one

To a solution of 12 g (0.070 mol) of 3,4-dihydrobenzo[b]furo[2,3-c]pyridine dissolved in 500 mL of acetonitrile at 60°C was added 20 g (0.140 mol) of 2-trimethylsiloxy-1,3-butadiene followed by 13.6 g (0.10 mol) of anhydrous zinc chloride. The mixture was heated at 60°C for 1.5 hour, cooled to 25°C, and 30 mL of 5% HCl was added and stirred 10 minutes. 40% Sodium hydroxide was added until the reaction was basic; 200 mL of water was added; and the acetonitrile layer was separated. The aqueous layer was filtered through celite and washed with ether. The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to a brown residue which was chromatographed (silica, ethyl acetate/hexane (1:1)) to give 8.2 g of product, m.p. 108-9°C.

Resolution of (12bSR)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3a]quinolizin-2-one

A solution of (-)-di-p-toluoyl-L-tartaric acid monohydrate (25.9 g) in 100 ml of ethyl acetate was mixed with a solution of (12bSR)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-one (15.5 g) in 700 ml of ethyl acetate and allowed to stand 12-78 hours. The mixture was filtered to yield 21 g of the di-p-toluoyl-L-tartrate salt of the amine. The free base was liberated by partitioning between aqueous Na<sub>2</sub>CO<sub>3</sub> and ethyl acetate ([α]<sub>D</sub> = ca. -79°; C=1; CHCl<sub>3</sub>). The diastereomeric salt of this material was again prepared following the above procedure. The collected di-p-toluoyl-L-tartrate salt was

2291S/0976A


- 17 -

17226IA

partitioned between ethyl acetate and aqueous  $\text{Na}_2\text{CO}_3$ , dried ( $\text{Na}_2\text{SO}_4$ ), filtered, treated with charcoal, filtered and evaporated to yield 5.4 g (35%) of (12bS)-1,3,4,6,7,12b-hexahydrobenzo[b]furo-  
 5 [2,3-a]quinolizin-2-one;  $[\alpha]_D = -84^\circ$ ; (C=1,  $\text{CHCl}_3$ ).

The (12bR)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-one was obtained by substituting (+)-di-p-toluoyl-D-tartaric acid monohydrate for (-)-di-p-toluoyl-L-tartaric acid in  
 10 the above procedure to provide product with  $[\alpha]_D = +84^\circ$  (C=1,  $\text{CHCl}_3$ ).

Employing the procedures substantially as described in Example 1, Steps A through E/F, or in some cases, Steps C through E/F but substituting for  
 15 the 3-benzofuranone used in Step A thereof the ketones described in Table I, or for the ethylamines used in Step C thereof, the corresponding ethylamines described in Table I, or for the butadienes used in  
 Step E thereof, the corresponding substituted  
 20 butadienes described in Table I, there are prepared the Ar[2,3-a]quinolizin-2-ones, also described in Table I by the following reactions:



2291S/0976A

- 18 -

17226IA

TABLE I

	<u>Ar</u>	<u>R<sup>5</sup></u>	<u>Ar</u>	<u>R<sup>5</sup></u>
5	10-chlorobenzo-	H	benzo[b]thieno-	1-CH <sub>3</sub>
	[b]furo-	H	10-methylbenzo-	4-COOCH <sub>3</sub>
	thieno-		[b]thieno-	
10	furo-	3-CH <sub>3</sub>	9-methoxybenzo-	H
			[b]thieno-	
	11-hydroxy-	H	11-fluorobenzo-	H
	benzo[b]furo		[b]furo-	
	10,11-dimethyl-	H	9-bromobenzo-	1-CON(CH <sub>3</sub> ) <sub>2</sub>
15	benzo[b]furo-		[b]furo-	
			11-methoxybenzo-	H
			[b]furo-	
	pyridino-	H	thiazolo-	H
	imidazo	H	pyrazolo-	H

20

Step F: Preparation of (2RS,12bSR)-2-aminomethyl-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine

Into a 1000 ml flask was placed 7 g (29 mmol) of 1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-one in 500 ml dry THF which had previously been saturated with dry methylamine at 0°C. To this mixture was added 10.44 g (64 mmol) of diethyl cyanophosphonate. After stirring for 18 hours, the solvent was removed and the resultant crude aminonitrile was dissolved in 300 ml dry THF and treated with 145 ml 1M borane in THF. This mixture was refluxed for 18 hours, cooled and quenched by the slow addition of methanol until

2291S/0976A

- 19 -

172261A

ebullition ceased, after which 400 ml 6N HCl was added and the reaction mixture was refluxed for an additional 2 hours. After cooling, the solvent was removed and the residue was basified by the addition  
5 of 400 ml saturated  $\text{Na}_2\text{CO}_3$  solution. This was extracted with 5 X 100 ml  $\text{CHCl}_3$ . The combined organic extracts were dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent was evaporated. Medium pressure column chromatography (chloroform saturated with ammonia)  
10 yielded 0.552 g. (7%) (2SR,12bSR)-2-aminomethyl-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine as a yellow oil, followed by 4.9 g of the desired product (62%) (2RS, 12bSR)-2-aminomethyl-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]  
15 quinolizine, also as a yellow oil.

Similarly prepared are the (2R,12bs)- and (2S,12bR) enantiomers by starting with the enantiomeric quinolizin-2-ones described in Step E.

20 Step G: Preparation of (2RS,12bSR)-3'-methyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo-[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one

Into a 400 ml flask was placed 4.4 g (15.8 mmol) of (2RS,12bSR)-2-aminomethyl-2-methylamino-1,3,  
25 4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]-quinolizine from Step F in 200 ml toluene. To this was added 5 g (32 mmol) of 1,1'-carbonyldiimidazole and the reaction was stirred for 5 hours, after which the toluene was washed with 3 X 50 ml  $\text{H}_2\text{O}$ , 50 ml brine,  
30 dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent evaporated to obtain a yellow solid. This material was dissolved in hot ethyl acetate, decolorized, filtered and treated with ethanolic HCl to give 3.53 g (72%) of (2RS,12bSR)-3'-methyl-spiro(1,3,4,6,7,12b-hexa-

2291S/1163A


- 20 -

17226IA

hydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one as a white crystalline hydrochloride salt which was recrystallized from methanol/ethyl acetate: m.p. 220°C (dec).

5 Similarly prepared are the (2R,12bs)- and (2S,12bR)-enantiomers of the 3'-methylimidazolidin-2'-one by starting with the enantiomeric diamines described in Step F hereof.

Employing the procedure substantially as  
10 described in Example 1, Steps F and G but substituting for the 2-aminomethyl-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine and methylamine used in Step F thereof an equimolar amount of the quinolizin-2-ones described in Table I and  
15  $R^3NH_2$ , there are prepared the spiro-imidazolidin-2-ones described in Table II in accordance with the following reaction scheme:



2291S/0976A

- 21 -

172261A

TABLE II

5	<div>Ar</div>	$R^5$		<div>Ar</div>	$R^5$	$R^3$
	10-chlorobenzo- [b]furo-	H	-CH <sub>3</sub>	benzo[b]thieno-	1-CH <sub>3</sub>	-CH <sub>3</sub>
10	thieno	H	H	10-methylbenzo- [b]thieno-	4-COOCH <sub>3</sub>	-CH <sub>3</sub>
	furo	3-CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	9-methoxybenzo- [b]thieno-	H	H
15	11-hydroxy- benzo[b]furo	H	-C <sub>3</sub> H <sub>7</sub>	11-fluorobenzo- [b]furo-	H	H
20	10,11-dimethyl- benzo[b]furo-	H	-CH <sub>3</sub>	9-bromobenzo- [b]furo-	1-CON(CH <sub>3</sub> ) <sub>2</sub>	PhCH <sub>2</sub> -
				11-methoxybenzo- [b]furo-	H	H
25	pyridino-	H	-CH <sub>3</sub>	thiazolo-	H	CH <sub>3</sub>
	imidazo	H	-CH <sub>3</sub>	pyrazolo-	H	CH <sub>3</sub>

30

2291S/0976A

- 22 -

17226IA

EXAMPLE 2

(2RS,12bSR)-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo  
[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one

5      Step A: Preparation of (2RS,12bSR)-2-aminomethyl-  
2-amino-1,3,4,6,7,12b-hexahydrobenzo[b]-  
furo[2,3-a]quinolizine

10      Employing the procedure substantially as  
described in Example 1, Step F, but substituting for  
the methylamine used therein, an equimolecular amount  
of ammonia, there was produced the title compound in  
comparable yield which was used directly in the next  
step without characterization.

15      Step B: Preparation of (2RS,12bSR)-spiro(1,3,4,6,7,  
12b-hexahydro[b]furo[2,3-a]quinolizin)-2,4'-  
imidazolidin-2'-one

20      The mixture of 540 mg of the diamine from  
Step A and 648 mg of 1,1'-carbonyldiimidazole in 200 ml  
of toluene was stirred overnight under N<sub>2</sub>. To this  
reaction mixture, 10 ml of H<sub>2</sub>O was added and  
stirred for 10 minutes. A precipitated solid was  
collected by filtration and transformed into HCl  
salt. The salt was recrystallized from MeOH-CHCl<sub>3</sub>,  
yield 80 mg, m.p. 270°C.

25      Employing the procedure substantially as  
described in Example 1, but substituting for the  
methylamine used therein an equimolecular amount of  
n-propylamine, there are produced in sequence:

30      (2RS,12bSR)-2-aminomethyl-2-n-propylamino-1,3,4,6,  
7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine, and

(2RS,12bSR)-3'-n-propyl-spiro(1,3,4,6,7,12b-hexahydro-  
benzo[b]furo[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one.

2291S/0976A

- 23 -

172261A

EXAMPLE 3

(2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one

---

5                    Into a 125 ml Erlenmeyer flask was placed  
0.225 g (0.72 mmol) (2SR,12bRS)-3'-methyl-spiro(1,3,4,  
6,7,12b-hexahydrobenzo[b]furo-[2,3-a]quinolizin)-2,4'-  
imidazolidin-2'-one (Example 1) in 50 ml toluene  
followed by 50 ml 40% NaOH, 0.366 g (1.08 mmol)  
10    tetrabutylammonium hydrogen sulfate and, with  
vigorous stirring, 0.067 ml (1.08 mmol) methyl  
iodide. Stirring was continued for 2 hours, after  
which the toluene layer was separated and washed with  
3 X 50 ml H<sub>2</sub>O, 50 ml brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and  
15    the solvent was removed to obtain 0.2 g (85%) pure  
(2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-  
hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-  
imidazolin-2'-one. This was dissolved in 50 ml ethyl  
acetate and ethanolic HCl was added to give a white  
20    crystalline hydrochloride salt: m.p. 260°C (dec.).

Employing the above procedure but starting  
with the enantiomers of the 3'-methyylimidazolidin-2'-  
one described in Example 1, Step G, there were  
produced the (2R,12bS)- and (2S,12bR)-enantiomers of  
25    the title compound with  $[\alpha]_D^{+1^\circ}$  (C=1.0 CH<sub>3</sub>OH),  
and  $[\alpha]_D^{-1^\circ}$  (C=1.0 CH<sub>3</sub>OH) respectively.

EXAMPLE 4

(2RS,12bSR)-1'-ethyl-3'-methyl-spiro[1,3,4,6,7,12b-  
30    hexahydrobenzo[b]furo[2,3-a]quinolizin]-2,4'-  
imidazolidin-2'-one-HCl 0.25 H<sub>2</sub>O

---

To a solution of the 3'-methyylimidazoline  
compound from Example 1 (0.1 g, 0.32 mmol) in 20 ml



2291S/1163A

- 24 -

172261A

of toluene was added tetrabutylammonium hydrogen sulfate (0.163 g, 0.48 mmol), 20 mls of 40% sodium hydroxide solution, and, with vigorous stirring, ethyl iodide (0.075 g, 0.48 mmol). This was stirred  
5 for 18 hours and then the toluene layer was separated and washed with 3 X 30 ml of water, 30 ml of brine, dried ( $\text{Na}_2\text{SO}_4$ ) filtered and concentrated to give an oil which was chromatographed (silica, ethyl acetate) to give the product as the  $\text{HCl}$   $0.25 \text{ H}_2\text{O}$ ,  
10 m.p. 235-239°C.

#### EXAMPLE 5

(2RS,12bSR)-3'-methyl-1'-propyl-spiro[2,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-  
15 imidazolidin-2'-one HCl

To a solution of the 3'-methylimidazoline compound from Example 1 (0.1 g, 0.32 mmol) in 20 ml of toluene was added tetrabutylammonium hydrogen sulfate (0.163 g, 0.48 mmol), 20 mls of 40% sodium hydroxide  
20 solution, and, with vigorous stirring, N-propyl iodide (0.110 g, 0.64 mm). This was stirred for 5 hours and then the toluene layer was separated and washed with 3 X 30 ml of water, 30 ml of brine, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to give an oil which was  
25 chromatographed (silica, ethyl acetate) to give the product as the  $\text{HCl}$ ; m.p. 240-242°C.

#### EXAMPLE 6

(2RS,12bSR)-1'-hydroxyethyl-3'-methyl-spiro[1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin]-2,4'-  
30 imidazolidin-2'-one HCl,  $0.25 \text{ H}_2\text{O}$

Step A: Preparation of 2-methoxy-2-(2-iodoethoxy) propane

To 6 mls of cold (0°C) methoxypropene was added 3 mls of 2-iodoethanol and 1 drop of phosphorous

2291S/0976A

- 25 -

172261A

oxychloride ( $\text{POCl}_3$ ). The reaction was stirred for 1 hour and then solid potassium carbonate was added. After 10 minutes the liquid was decanted and concentrated to give the product as an oil.

5

Step B: (2RS,12bSR)-1'-hydroxyethyl-3'-methyl-spiro [1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin]-2,4'-imidazolidin-2'-one HCl, 0.25  $\text{H}_2\text{O}$

10

To a solution of the 3'-methylimidazolidine from Example 1 (0.04 g, 0.16 mmol) in 20 ml of toluene was added tetrabutylammonium hydrogen sulfate (0.082 g, 0.24 mmol), 20 mls of 40% sodium hydroxide solution, and, with vigorous stirring, 2-methoxy-2-(2-iodoethoxy)propane (0.053 g, 0.209 mmol). This was stirred for 45 hours and then the toluene layer was poured into 20 mls of 5% HCl solution, stirred for 15 minutes and then made basic. The toluene layer was separated and washed with 3 X 30 ml of water, 30 ml of brine, dried ( $\text{Na}_2\text{SO}_4$ ) filtered and concentrated to give an oil which was chromatographed (silica,  $\text{NH}_3$ /saturated  $\text{CHCl}_3$ ) to give the product as the HCl, (0.25  $\text{H}_2\text{O}$ ; m.p. 172-176°C.

25

#### EXAMPLE 7

(2RS,12bSR)-1'-Benzyl-3'-methyl-spiro[1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin]-2,4'-imidazolidin-2'-one HCl, 0.25  $\text{H}_2\text{O}$

30

To a solution of the 3'-methylimidazolidine compound from Example 1, (0.1 g, 0.32 mmol) in 20 ml of toluene was added tetrabutylammonium hydrogen sulfate (0.163 g, 0.48 mmol), 20 mls of 40% sodium

2291S/1163A

- 26 -

17226IA

hydroxide solution, and, with vigorous stirring, benzyl bromide (0.123 g, 0.720 mmol). This was stirred for 5 hours and then the toluene layer was separated and washed with 3 X 30 ml of water, 30 ml of  
5 brine, dried ( $\text{Na}_2\text{SO}_4$ ) filtered and concentrated to give an oil which was chromatographed (silica, ethyl acetate) to give the product as the HCl, 0.25  $\text{H}_2\text{O}$ ; m.p. 245-248°C.

10

EXAMPLE 8

(2RS,12bSR)-1'-acetyl-3'-methyl-spiro[1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin]-2,4'-imidazolidin-2'-one HCl, 0.25  $\text{H}_2\text{O}$

To a solution of the 3'-methyylimidazolidine  
15 from Example 1 (0.255 g, 0.5 mmol) in 15 ml of THF (0°C) was added 0.345 mls of a 1.6 M n-butyl lithium solution (0.55 mmol). After 15 minutes the reaction was cooled to -78°C and acetyl chloride (0.050 g, 0.64 mmol) was rapidly added. Stirring was continued  
20 at -78°C for 30 minutes and the temperature was then raised to 25°C over 2 hours. The reaction mixture was then poured into 30 ml of water, extracted with 2 x 20 ml of ethyl acetate, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to an oil which was  
25 chromatographed (silica, ethyl acetate) to give the product as the HCl, 0.25  $\text{H}_2\text{O}$ ; 251-255°C.

EXAMPLE 9

(2RS,12bSR)-1'-acetyl-3'-propyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one

Employing the procedure substantially as described in Example 8, but substituting for the

2291S/1163A

- 27 -

17226IA

3'-methylimidazoline compound used in Example 8, an equimolecular amount of the corresponding 3'-propyl-imidazolidine compound, there is obtained, the title compound in comparable yield.

5

EXAMPLE 10

(2RS,12bSR)-1'-methyl-3'-propyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one

---

10           Employing the procedure substantially as described in Example 3, but substituting for the 3'-methylimidzolidine compound used in Example 3, an equimolecular amount of the corresponding 3'-propylimidazoline compound, there is obtained, the  
15   title compound in comparable yield; m.p. 220-225°C.

EXAMPLE 11

(2RS,12bSR)-3'-methyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-imidazolidin-2'-thione

---

20           (2RS,12bSR)-2-Aminomethyl-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine (285 mg, 1 mmole) and 1,1'-thiocarbonyldiimidazole (356 mg, 2 mmole) were stirred in toluene (20 ml) at  
25   ambient temperature for 2 hours then at 80°C for 2 hours. The reaction mixture was concentrated to dryness, dissolved in CH<sub>2</sub>Cl<sub>2</sub>, washed with H<sub>2</sub>O, saturated NaCl solution, and dried (Na<sub>2</sub>SO<sub>4</sub>). Filtration and concentration to dryness provided 88  
30   mg (27%) of the subject compound after chromatography. Treatment with ethanolic HCl provided the hydrochloride salt; m.p. 280°C.

2291S/0976A

- 28 -

172261A

Anal. for  $C_{18}H_{21}N_3OS \cdot HCl \cdot 1/2 H_2O$ :  
calc'd.: C, 57.98; H, 6.22; N, 11.27.  
Found: C, 58.15; H, 6.23; N, 11.10.

5

EXAMPLE 12

(2RS,12bSR)-2'-methyl-spiro(1,3,4,6,7,12b-hexahydro-benzo[b]furo-[2,3-a]quinolizin)-2,3'-(1,2,5-thiadiazolidin-1',1'-dioxide), HCl

---

- 10 A mixture of 570 mg of the diamine from Example 1, Step I, and 1 ml of triethylamine in 20 ml of  $CHCl_3$  was stirred at  $-78^\circ C$  under  $N_2$ . Sulfuryl chloride (323 mg) in 10 ml of  $CHCl_3$  was added dropwise. The mixture was stirred at  $-78^\circ C$  for 1 hour and at room temperature overnight. Evaporation
- 15 gave an oily residue which was redissolved in  $CHCl_3$ , washed with  $H_2O$ , dried over  $K_2CO_3$ , and evaporated to dryness. The product was purified by spinning disk chromatography and transformed into the HCl salt with ethanolic hydrogen chloride.
- 20 Recrystallization from ethanol-ether gave 15 mg of product; m.p.  $250^\circ C$ .

EXAMPLE 13

25 (2RS,12bSR)-3'-Methyl-spiro(1,3,4,6,7,12b-hexahydro-benzo[b]thieno[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one

---

- 30 To a three necked, oven dried round bottomed flask with stirring bar was added 100 ml of dry THF. The flask and its contents were cooled in an ice bath to  $0^\circ C$  and the THF was saturated with methylamine gas. To this solution was added 590 mg of 1,3,4,6,7,12b-hexahydrobenzo[b]thieno[2,3-a]quinolizin-2-one and 816 mg of diethylcyanophosphate under the

2291S/0976A

- 29 -

17226IA

protection of a calcium sulfate drying tube. The ice bath was allowed to expire and the mixture was stirred overnight at room temperature. The reaction mixture was concentrated in vacuo, and redissolved in 50 ml of dry THF to which was added 25 ml of 1.0M borane-THF complex, under argon. This mixture was heated at reflux overnight. The cooled reaction mixture was quenched with 3 ml of methanol, then 20 ml of 2N HCl were added and the THF was removed in vacuo. The aqueous residue was made basic with concentrated ammonium hydroxide and extracted with chloroform (2 x 25 ml). The combined chloroform extracts were dried over anhydrous potassium carbonate, filtered and concentrated in vacuo. Chromatography of the crude product on 20 g of silica gel with 2.5% methanol in ammonia saturated chloroform as eluant provided 170 mg of intermediate diamine.

To a 100 ml round bottomed flask containing 170 mg of the aforementioned diamine was added 3 ml of chloroform and 15 ml of toluene. To this solution was added 406 mg of carbonyldiimidazole and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with 100 ml of ethyl acetate and was washed with water and brine. Drying (potassium carbonate), filtration and removal of the solvent in vacuo left an off-white solid. Chromatography of this material on 20 g of silica gel with ammonia saturated chloroform provided 83 mg of the title compound as a crystalline solid. An analytical sample was prepared by recrystallization from ethanol-ethyl acetate; m.p. 290°C (dec.).

2291S/1163A

- 30 -

17226IA

EXAMPLE 14

(2RS,12bSR)-1',3'-Dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]thieno[2,3-a]quinolizine)-2,4'-imidazolidin-2'-one

5           To a 50 ml round bottomed flask was added  
(2RS,12bSR)-3'-methyl-spiro(1,3,4,6,7,12b-hexahydro-  
benzo[b]thieno[2,3-a]quinolizine)-2,4'-imidazolidin-  
2'-one and 10 ml of toluene. When the material had  
dissolved, 10 ml of 10N sodium hydroxide and 44 mg of  
10   tetrabutylammonium sulfate were added. While  
stirring vigorously, 8.1 microliters of methyl iodide  
was added.

The mixture was stirred for 30 minutes at  
room temperature. The layers were separated and the  
15   toluene layer was washed with water and brine.  
Drying (potassium carbonate), filtration and removal  
of the solvent in vacuo left 29 mg of crude product.  
This material was chromatographed on 3 g of silica  
gel with 3% (V/V) methanol in chloroform. The oil  
20   thus obtained was triturated with ethyl acetate and  
the crystals that precipitated were collected on a  
frit and dried overnight at 70° C/0.05 torr giving 25  
mg of the title compound; m.p. 204-206°C.

25

EXAMPLE 15

(2RS,12bSR)-3'-Methyl-spiro(10-methyl-1,3,7,12b-hexahydrobenzo[b]thieno[2,3-a]quinolizine)-2,4'-imidazolidin-2'-one

30           To an oven dried 100 ml round bottomed flask  
with stirring bar, argon inlet and a gas dispersion  
tube was added 75 ml of dry THF. The contents of the  
flask were cooled to 0°C and saturated with  
methylamine gas. To this solution was added 400 mg

of 10-methyl-1,3,4,6,7,12b-hexahydrobenzo[b]thieno  
[2,3-a]quinolizin-2-one and 0.491 ml of diethylcyano-  
phosphonate. The ice bath was allowed to expire and  
the mixture was stirred at room temperature  
5 overnight. The entire mixture was concentrated  
in vacuo and redissolved in THF. To this solution  
was added 10.2 ml of a 0.98M solution of borane-THF  
complex with a syringe. This mixture was heated at  
reflux over night. The cooled reaction mixture was  
10 quenched with methanol and concentrated to a  
colorless syrup in vacuo. This material was stirred  
with 100 ml of 2N HCl over night. The HCl solution  
was washed with ether, made basic with 20% NaOH, and  
extracted with chloroform (2 x 75 ml). The combined  
15 chloroform extracts were washed with brine and dried  
over anhydrous potassium carbonate. Filtration and  
removal of the solvent left 231 mg of a colorless  
oil. This material was chromatographed on 30 g of  
silica gel with 5% methanol in ammonia saturated  
20 chloroform as eluant. This chromatography provided  
51 mg of the intermediate diamine.

To a 100 ml round bottomed flask containing  
the aforementioned diamine (51 mg) was added 1.0 ml  
of dry chloroform. When the material had dissolved,  
25 5 ml of toluene was added followed by 81 mg of  
carbonyl diimidazole. This mixture was stirred  
overnight at room temperature under argon. The  
reaction mixture was diluted with 30 ml of ethyl  
acetate and washed with water (2 x 15 ml) and brine  
30 (25 ml). Drying over magnesium sulfate, filtration  
and removal of the solvent in vacuo left 60 mg of a  
solid which was chromatographed on 5 g of silica gel  
with ammonia saturated chloroform to give 28.6 mg of



2291S/0976A

- 32 -

17226IA

the title compound which crystallized on trituration with ethyl acetate; m.p. 225°C (dec.)

EXAMPLE 16

5 (2RS,12bSR)-3'-Methyl-spiro(11-chloro-1,3,4,6,7,12b-hexahydrobenzo[b]thieno[2,3-a]quinolizine)-2,4'-imidazolidin-2'-one

Step A: Preparation of (2RS,12bSR)-2-Aminomethyl-2-methylamino-11-chloro-1,3,4,5,7,12b-hexahydro-  
10 benzo[b]thieno[2,3-a]quinolizine

11-chloro-1,3,4,6,7,12b-hexahydrobenzo[b]thieno[2,3-a]quinolizine-2-one (500 mg, 1.71 mmole) was added to dry tetrahydrofuran (100 ml) saturated with methylamine with stirring at 0°C. Diethyl  
15 cyanophosphonate (0.7 ml, 4.6 mmole) was added; the solution was stoppered and left to stir at ambient temperature for 18 hours. The solvent was removed and the residue partitioned between ethyl acetate and H<sub>2</sub>O. The organic layer was separated, washed with  
20 H<sub>2</sub>O, saturated NaCl solution, and dried (Na<sub>2</sub>SO<sub>4</sub>). Filtration and concentration provided the aminonitrile which was immediately dissolved in fresh tetrahydrofuran (100 ml) and treated with borane in tetrahydrofuran (0.98 M, 10.42 ml, 10.4  
25 mmole). After stirring at ambient temperature for 15 minutes the reaction mixture was heated at reflux for 18 hours. After cooling, methanol was added slowly to destroy excess borane, 6N HCl (80 ml) was added and the mixture was heated at reflux for 2 hours.  
30 After cooling the reaction mixture was basified with solid NaOH, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to give 310 mg (54%) of title compound after chromatography.

2291S/0976A

- 33 -

17226IA

Step B: Preparation of (2RS,12bSR)-3'-Methyl-spiro  
(11-chloro-2,3,4,6,7,12b-hexahydro-2H-benzo-  
thieno[2,3-a]quinolizine)-2,4'-imidazolidin-  
2'-one

---

5 (2RS,12bSR)-2-Aminomethyl-2-methylamino-11-  
chloro-1,3,4,6,7,12b-hexahydrobenzo[b]thieno[2,3-a]  
quinolizine (270 mg, 0.8 mmole) and 1,1'-carbonyl-  
diimidazole (254 mg, 1.6 mmole) were dissolved in  
toluene (100 ml) and stirred at ambient temperature  
10 under N<sub>2</sub> for 5 hours. Water was added; the organic  
layer was separated, washed with H<sub>2</sub>O, saturated  
NaCl solution and dried (Na<sub>2</sub>SO<sub>4</sub>). Filtration and  
concentration followed by chromatography provided 140  
mg (42%) of the subject compound. Treatment with  
15 ethanolic HCl gave the hydrochloride salt, m.p.  
322-325°C.

Anal. for C<sub>17</sub>H<sub>20</sub>ClN<sub>3</sub>OS.HCl.H<sub>2</sub>O:

Calc'd: C, 51.92; H, 5.58; N, 10.09.

Found: C, 51.94; H, 5.34; N, 9.80.

20

#### EXAMPLE 17

(2RS,12bSR)-spiro[1,3,4,6,7,12b-hexahydrobenzo[b]furo  
[2,3-a]quinolizin]2,5'-tetrahydrofuran-2'-one HCl,  
0.25 H<sub>2</sub>O

---

25 To 2.85 ml of 2.6 M n-butyllithium (4.0  
mmol) at -40°C was added a solution of allyl-bis-  
dimethylamino phosphonate in 0.5 ml of THF. The  
temperature was raised to -20°C and stirring was  
continued for 2 hours, after which a solution of  
30 1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-  
2-one in 1 ml of THF was added. The reaction was  
allowed to come to room temperature, stirred for 2.5  
hours and was poured into 50 mls of water which was  
extracted with 2 x 30 ml of ethyl acetate, dried

2291S/0976A

- 34 -

17226IA

(Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to give a residue which was chromatographed (silica, ethyl acetate) to give the product. Treatment with ethanol-HCl gave the HCl 0.25 H<sub>2</sub>O salt, m.p. 258-262°C.

#### EXAMPLE 18

(2SR,12bSR)-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,5'-oxazolidin-2'-one

10 Step A: Preparation of (2SR,12bSR)-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,2'-oxirane

In a 50 ml 3 neck flask 0.053 g (1.1 mmol) sodium hydride as a 50% oil dispersion was washed three times with toluene under nitrogen. The resulting solid was then slurried in 10 ml dry DMF and cooled to 0°C. To this was added 0.25 g (1.1 mmol) trimethylsulfoxonium iodide. After stirring for 20 minutes, 0.241 g (1 mmol) of 1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-one in 4 ml dry DMF was added dropwise. Stirring was continued for 10 minutes, after which the reaction was poured into 100 ml H<sub>2</sub>O and extracted with 4 x 20 ml ethyl acetate. The combined organic extracts were washed with 25 ml brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent was removed to obtain 0.25 g (98%) of 2SR,12bSR-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,2'-oxirane as a yellow solid which was recrystallized from ether/pet. ether; m.p. 109-110°C.

30

2291S/0976A

- 35 -

17226IA

Step B: Preparation of (2SR,12bSR)-2-aminomethyl-2-hydroxy-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine

Into a pressure bottle was placed 0.19 (0.4 mmol) (2SR,12bSR)-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,2'-oxirane dissolved in 10 ml absolute ethanol. After cooling the vessel to -78°C using a dry ice/acetone bath, 10 ml ammonia was condensed into the bottle. The bottle was sealed, warmed to room temperature and allowed to stir for 48 hours after which the pressure was released and the solvent removed in vacuo. Subsequent spinning disc chromatography (5% methanol/chloroform saturated with ammonia) yielded .071 g. (83%) of (2SR,12bSR)-2-aminomethyl-2-hydroxy-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine which was recrystallized from hexane/ethyl acetate; m.p. 155-157°C.

Step C: Preparation of (2SR,12bSR)-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,5'-oxazolidin-2'-one

In a 50 ml flask, 0.093 g (0.34 mmol) (2SR,12bSR)-2-aminomethyl-2-hydroxy-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine in 20 ml toluene was placed, and to this was added 0.102 g (0.62 mmol) 1,1'-carbonyldiimidazole. After refluxing for 1 hour, the reaction was cooled and washed with 3 x 10 ml H<sub>2</sub>O, 10 ml brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed to give 0.096 g (95%) (2SR,12bSR)-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,5'-oxazolidin-2'-one which was dissolved in 50 ml ethyl acetate. Ethanolic

2291S/1163A

- 36 -

17226IA

HCl was then added dropwise to give a white hydrochloride hemihydrate salt which was recrystallized from ethyl acetate, m.p. 280°C (dec.).

5

EXAMPLE 19

(2RS,12bSR)-3'-Methyl-spiro(1,3,4,6,7,12b-hexahydro-benzo[b]furo[2,3-a]quinolizine)-2,4'-oxazolidin-2'-one

10

Step A: Preparation of (2RS,12bSR)-2-Cyano-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine

A mixture of 1,3,4,6,7,12b-hexahydro-benzo[b]furo[2,3-a]quinolizine-2-one (2.4 g, 9.9 mmole) and MgSO<sub>4</sub> (250 mg) in dry tetrahydrofuran (600 ml) was saturated with methylamine at 0°C. Diethylcyanophosphonate (3.35 ml, 22.1 mmole) was added in one portion, and the mixture was left stoppered for 5 days. The reaction mixture was filtered; concentrated to dryness and the residue partitioned between ethyl acetate and H<sub>2</sub>O; the organic layer separated; washed with H<sub>2</sub>O, saturated NaCl solution; and dried (Na<sub>2</sub>SO<sub>4</sub>). Filtration and concentration gave crude subject compound (4.0 g) which was used immediately in the next step.

25

Step B: Preparation of (2RS,12bSR)-Ethyl-2-methylamino-1,3,4,6,7,12b-hexahydro-benzo[b]furo[2,3-a]quinolizine-2-carboxylate

30

Sulfuric acid (36N) (30 ml) was added to a solution of (2RS,12bSR)-2-cyano-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine (3.2 g, 0.011 mole) in 95% ethanol (30 ml) with cooling at 0°C. After heating at reflux for 3 hours,

2291S/0976A

- 37 -

17226IA

the reaction mixture was cooled to 0°C, treated with ice-H<sub>2</sub>O, basified with 20% NaOH solution and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, washed with saturated NaCl solution and  
5 dried (Na<sub>2</sub>SO<sub>4</sub>). Filtration and concentration followed by flash chromatography provided 930 mg (26%) of the subject compound.

Step C: Preparation of (2RS,12bSR)-2-Hydroxy-  
10 methyl-2-methylamino-1,3,4,6,7,12b-hexa-  
hydrobenzo[b]furo[2,3-a]quinolizine  
(2RS, 12bSR)-Ethyl-2-methylamino-  
1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-  
carboxylate (940 mg, 2.86 mmole) in dry diethyl ether  
15 (20 ml) was added dropwise to a refluxing suspension of lithium aluminum hydride (108 mg, 2.86 mmole) in diethyl ether (20 ml) with stirring under N<sub>2</sub>. After heating at reflux for 2 hours, the reaction mixture was cooled to 0°C, the excess hydride was  
20 destroyed, and the mixture was left to stir for 16 hours. Filtration, followed by extraction of the salts with CH<sub>2</sub>Cl<sub>2</sub> provided 560 mg (68%) of the title compound after chromatography.

25 Step D: Preparation of (2RS,12bSR)-3'-Methyl-spiro  
(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]  
quinolizine)-2,4'-oxazolidin-2'-one  
(2RS, 12bSR)-2-Hydroxymethyl-2-methylamino-  
1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo[2,3-a]-  
30 quinolizine (200 mg, 0.7 mmole) and 1,1'-carbonyl-diimidazole (222 mg, 1.37 mmole) in toluene (120 ml) were stirred at ambient temperature for 18 hours. Water was added, the organic layer separated,

2291S/0976A

- 38 -

172261A

extracted with saturated NaCl solution, dried  
(Na<sub>2</sub>SO<sub>4</sub>), then filtered and concentrated to give  
the title compound. Treatment with ethanolic HCl  
provided the hydrochloride salt (184 mg, 86.7%), m.p.  
5 265-267°C.

Anal. for C<sub>18</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub>.HCl 0.5 H<sub>2</sub>O:

Calc'd: C, 60.41; H, 6.21; N, 7.83.

Found: C, 60.42; H, 6.05; N, 7.72.

10

EXAMPLE 20

(2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-  
hexahydrobenzo[b]furo[2,3-a]quinolizine)-2,4'-  
imidazolidin-2',5'-dione

Methyl isocyanate (19 mg, 19.8 µl, 0.34  
15 mmole) was added to a solution of (2RS, 12bSR)-  
ethyl 2-methylamino-1,3,4,6,7,12b-hexahydro-2H-  
benzo[b]furo[2,3-a]quinolizine-2-carboxylate (100 mg,  
0.30 mmole) in acetonitrile (10 ml) at ambient  
temperature with magnetic stirring under N<sub>2</sub>. After  
20 stirring for 18 hours the acetonitrile was removed,  
and the residue was dissolved in CHCl<sub>3</sub>, washed with  
H<sub>2</sub>O, saturated NaCl solution, and dried  
(Na<sub>2</sub>SO<sub>4</sub>). Filtration and concentration to  
dryness followed by conversion to the hydrochloride  
25 salt with ethanolic HCl gave 100 mg (87.4%) of  
(2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexa-  
hydrobenzo[b]furo[2,3-a]quinolizine)-2,4'-imidazolin-  
2',5'-dione, m.p. 204-206°C.

30

2291S/0976A

- 39 -

172261A

EXAMPLE 21

(2RS,12bSR)-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]  
furo[2,3-a]quinolizin)-2,5'-oxazolidin-2'-one

5      Step A: Preparation of (2RS,12bSR)-spiro(1,3,4,6,7,  
12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-  
2,2'-oxirane

Into a mixture of n-butyllithium (1.6 M/  
hexane, 3.5 ml) and  $(CH_3)_2S^+CH_3I^-$  (1.23 g)  
in THF (25 ml), 1,3,4,6,7,12b-hexahydrobenzo[b]  
10 furo[2,3-a]quinolizin-2-one (400 mg) in THF (5 ml)  
was added at -5°C with stirring under N<sub>2</sub> gas. The  
mixture was stirred at 0°C for 30 minutes and then at  
room temperature overnight. To the mixture, were  
added ethyl acetate and water with vigorous stirring  
15 and then an insoluble solid was filtered off. The  
ethyl acetate layer was separated, washed with H<sub>2</sub>O  
and dried over MgSO<sub>4</sub>. Evaporation of the ethyl  
acetate gave a yellow oil which was subjected to  
silica-gel column chromatography to give the oxirane  
20 which was recrystallized from a mixture of ether-  
petroleum ether to give yellow prisms; yield 70 mg;  
m.p. 108-109°C.

25      Step B: Preparation of (2RS,12bSR)-2-aminomethyl-2-  
hydroxy-1,3,4,6,7,12b-hexahydrobenzo[b]furo  
[2,3-a]quinolizine

An alcohol solution (15 ml) of the oxirane  
(170 mg) was taken into a pressure bottle and cooled  
in dry ice-acetone. After adding liquid NH<sub>3</sub> (15  
30 ml) to the ethanol solution, the mixture was left for  
3 weeks with stirring. Evaporation of the ethanol  
gave a crude oil of the amino alcohol, (190 mg.)



2291S/1163A

- 40 -

172261A

Step C: Preparation of (2RS,12bSR)-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,5'-oxazolidin-2'-one

The amino alcohol from Step B (crude 180 mg) and 5 1,1'-carbonyldiimidazole (225 mg) were mixed in toluene (60 ml) under N<sub>2</sub> gas and stirred overnight at room temperature. Water (5 ml) was added to the reaction mixture and it was stirred for 20 minutes. The toluene layer was separated and the water layer 10 was washed with ethyl acetate. The organic layers were combined and washed with brine and water and dried over K<sub>2</sub>CO<sub>3</sub>. Evaporation gave a brown oil which was purified on chromatotron (a spinning thin layer chromatographic apparatus from Harrison 15 Research, California) to give the oxazolidone. The oxazolidone was transformed into the HCl salt and recrystallized from ethanol-ether; yield 20 mg; m.p. 270°C (dec.).

20

EXAMPLE 22

(2RS,12bSR)-3'-methyl-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2,5'-oxazolidin-2'-one

Employing the procedure substantially as described in Example 3, but substituting for the 25 (2SR,12bRS)-3'-methyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one used therein, an equimolar amount of the spiro-oxazolidin-2-one from Example 21, the subject compound is produced in comparable yield.

30

2291S/0976A

- 41 -

17226IA

EXAMPLE 23

(2RS,12bSR)-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]  
furo[2,3-a]quinolizine)-2,5'-pyrazolidin-3'-one

5     Step A: Preparation of (E,Z)-2-Carboethoxymethyl-  
          idine-1,3,4,6,7,12b-hexahydrobenzo[b]furo  
          [2,3-a]quinolizine

To a 3 necked 250 ml round bottomed flask  
with a stirring bar, argon inlet, thermometer and  
septum was added 8.11 g of a 28.6% suspension of  
10     potassium hydride in mineral oil. The oil was  
removed with two washings of hexane. To the oil free  
potassium hydride was added 25 ml of THF and the  
mixture was cooled to 0°C. A solution of triethyl-  
phosphonoacetate (12.98 g) in 15 ml of THF was added dropwise  
15     over 15 minutes. To this solution was added a  
solution of 3.00 g of 1,3,4,6,7,12b-hexahydrobenzo-  
[b]furo[2,3-a]quinolizine-2-one in 25 ml of dry THF,  
dropwise over 15 minutes. This mixture was allowed  
to stir overnight at room temperature. The mixture  
20     was diluted with 250 ml of ethyl acetate and was  
washed with water (3 x 200 ml) and brine (200 ml).  
Drying (magnesium sulfate), filtration and removal of  
the solvent in vacuo left an orange oil which was  
chromatographed on 300 g of silica gel with 30% ethyl  
25     acetate in hexanes as eluant. The chromatography  
provided two products: 650 mg of the faster eluting E  
isomer and 360 mg of the slower eluting Z isomer.

30     Step B: Preparation of (2RS,12bSR)-spiro-(1,3,4,6,7,  
          12b-hexahydrobenzo[b]furo[2,3-a]quinolizine-  
          2,5'-pyrazolidin-3'-one

To a 15 ml round bottomed flask with a  
stirring bar, reflux condenser, and argon inlet was

2291S/0976A

- 42 -

17226IA

added 600 mg of either of the above unsaturated esters and 5 ml of anhydrous hydrazine. Five milliliters of dry benzene were added to dissolve the ester. The mixture was heated at reflux in an oil bath for 1 hour. The reaction mixture was cooled to room temperature and the crystals which had precipitated were collected on a frit. The crystals were washed with benzene, dried in vacuo and recrystallized from boiling methanol to provide 251 mg of the title pyrazolidinone; m.p. 286-287°C.

#### EXAMPLE 24

(2RS,12bSR)-2'-methyl-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine)-2,5'-pyrazolidin-3'-one

Step A: Preparation of (E,Z)-2-Carbomethoxy-methylidene-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine

To 200 ml round bottomed flask with a mechanical stirrer was added 2.84 g of 35% suspension of potassium hydride in mineral oil, under argon. The oil was removed with two washings of hexanes and 15 ml of dry THF was added. The stirring suspension was cooled to 0°C and trimethylphosphonoacetate (4.53 g) was added, neat, dropwise. After this viscous mixture had stirred for 10 minutes a solution of 2.00 g of 1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine-2-one in 15 ml of dry THF was added with a syringe. The cooling bath was removed and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with 200 ml of water and extracted with ethyl acetate (3 x 100 ml). The combined ethyl acetate extracts were washed with

2291S/0976A

- 43 -

172261A

water and brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. The crude product was chromatographed on 150 g of silica gel with 25% ethyl acetate in hexanes as eluant. This procedure provided 1.25 g of the faster eluting E isomer and 1.12 g of the slower eluting Z isomer. The free bases were converted to their HCl salts by the usual method: E-isomer; m.p. 218-129°C; Z-isomer; m.p. 220-221°C.

The (12bR)- and (12bS)-enantiomers of the title compound were prepared by starting with the enantiomers of the quinolizin-2-one described in Example 1, Step F.

Step B: Preparation of (2RS,12bSR)-2'-Methyl-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine)-2,5'-pyrazolidin-3'-one dihydrochloride

To a 25 ml round bottomed flask with stirring bar, reflux condenser and argon inlet was added 228 mg of the E - $\alpha,\beta$ - unsaturated ester from step A, 10 ml of dry THF, and 407 microliters of methylhydrazine. This solution was heated at reflux for 36 hours. The cooled reaction mixture was concentrated in vacuo and the residue was crystallized from ethyl acetate-hexanes. Recrystallization of this material from ethyl acetate-hexanes gave 127 mg of the free base; m.p. 180-182°C. This material was converted into its dihydrochloride salt by dissolving it in boiling 2-propanol and adding two equivalents of ethanolic HCl. The crystals thus obtained were collected on a frit, washed with 2-propanol and dried to give 110 mg of the title compound; m.p. 270°C (dec.).

2291S/0976A

- 44 -

17226IA

EXAMPLE 25

(2RS,12bSR)-2'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-benzo[b]furo[2,3-a]-2,3'-isoxazolidin-5'-one

To a 25 ml round bottomed flask with  
5 stirring bar, argon inlet and a reflux condenser was  
added 264 mg of the E - $\alpha,\beta$ - unsaturated ester from  
step A of Example 24, 417 mg of N-methylhydroxylamine  
hydrochloride and 829 mg of finely powdered potassium  
10 carbonate. To this mixture was added 10 ml of dry  
THF with a syringe. This suspension was heated at  
reflux with vigorous stirring for 24 hours. The  
reaction mixture was diluted with 100 ml of ethyl  
acetate and washed with water and brine. Drying  
(potassium carbonate), filtration and removal of the  
15 solvent in vacuo gave a yellow crystalline solid.  
Recrystallization of this material from boiling ethyl  
acetate-hexanes gave 146 mg of the title compound;  
m.p. 183-185°C.

EXAMPLE 26

20 (2RS,12bSR)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-benzo[b]furo[2,3-a]quinolizine)-2,4'-azetid-2'-one

To an argon filled Parr hydrogenation bottle  
was added 500 mg of (2RS,12bSR)-2'-methyl-spiro-  
25 (1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]-2,3'-  
isoxazolidin-5'-one, 500 mg of 10% palladium on  
carbon, and 50 ml of methanol. This solution was  
hydrogenated on a Parr shaker at 50 psig and room  
temperature for 16 hours. The mixture was filtered  
30 through a celite pad and the catalyst was washed with  
a little methanol. The filtrates were combined and  
concentrated in vacuo. The residue was  
recrystallized from boiling 2-propanol and ethyl

2291S/1163A

- 45 -

172261A

ether to obtain 500 mg of intermediate beta-amino acid used in the following reaction.

To a 200 ml round bottomed flask with a stirring bar and argon inlet was added the  
5 aforementioned beta-amino acid, 50 ml of dry methylene chloride, 279 microliters of triethylamine, and 345 mg of ethyl-(3-dimethylaminopropyl)-carbodiimide hydrochloride. This solution was stirred under argon at room temperature for 70  
10 hours. The solution was diluted with methylene chloride and washed with water and brine. Drying (magnesium sulfate), filtration and removal of the solvent in vacuo gave 442 mg of a yellow oil. This oil was chromatographed on 30 g of silica gel with 3%  
15 methanol in chloroform as eluant. Trituration of the oil obtained from the chromatography with 1:1 ethyl acetate-hexanes gave 126 mg of light yellow crystals of the title compound; m.p. 173-175°C.

20

EXAMPLE 27

(2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-(5',6'-dihydro-1'H-pyrimidin-2'(3'H)-one, and  
(2SR,12bSR)-1',3'-dimethyl-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-(5',6'-  
25 dihydro-1'H-pyrimidin-2'(3'H)-one)

Step A: Preparation of (2RS,12bSR)-N-methyl-(2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide, and  
30 (2SR,12bSR)-N-methyl-(2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide

Into a pressure bottle was placed 2.8 g (9.4 mmol) (E,Z)-2-carbomethoxymethylidene-1,3,4,6,7,12b-

2291S/0976A

- 46 -

172261A

hexahydrobenzo[b]furo[2,3-a]quinolizine in 20 ml ethanol and the solution was cooled to -78°C with a dry ice/acetone bath. Methylamine (20 ml) was condensed into the vessel which was then sealed and allowed to stir at room temperature for 7 days. The pressure was released and the solvent removed in vacuo to yield 3.0 g (97%) (2RS,12bSR)-N-methyl-(2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide as a yellow oil.

The (12bR)- and (12bS)-enantiomers of the (2RS,12bSR)- title compounds were prepared by starting with the enantiomers of the carbomethoxymethylidene compound described in Example 24, Step A.

In a similar manner, 8.6 g (E,Z)-2-carbomethoxymethylidene-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine was reacted in a pressure vessel at 100°C for 18 hours in neat methylamine. Half of the crude reaction mixture (5 g) was subjected to medium pressure column chromatography (20% methanol/chloroform saturated with ammonia) to give 2.0 g (2RS,12bSR)-N-methyl-N-(2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl) acetamide as a yellow oil and 0.7 g of (2SR,12bSR)-N-methyl-N-(2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide.

Step B: Preparation of (2SR,12bSR)-2-methylamino-2-(2'-methylaminoethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine, and (2RS,12bSR)-2-methylamino-2-(2'-methylaminoethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine

Into a 500 ml flame dried flask was placed 10.8 ml (36.7 mmol) 3.4 M sodium bis (2-methoxyethoxy)

2291S/0976A

- 47 -

172261A

aluminum hydride in toluene and 150 ml dry THF.  
After heating to reflux, 3.0 gr. (9.1 mmol)  
(2RS,12bSR)-N-methyl-N-(2-methylamino-1,3,4,6,7,12b-  
hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide  
5 in 50 ml dry THF was added dropwise over a 10 minute  
period. Refluxing was continued for 3 hours after  
which time the reaction was cooled and quenched by  
the dropwise addition of a saturated sodium potassium  
tartrate solution. The solvent was removed and the  
10 residue partitioned between H<sub>2</sub>O/chloroform. The  
organic layer was separated, washed with 50 ml brine,  
dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent evaporated. Medium  
pressure column chromatography (10% methanol/chloro-  
form saturated with ammonia gave 1.6 g (56%)  
15 (2RS,12bSR)-2-methylamino-2-(2'-methylaminoethyl)-  
1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine  
as a yellow oil.

Similarly prepared are the enantiomers of  
the title compound from enantiomeric acetamides from  
20 Step A.

Similarly, 0.186 g (0.57 mmol) (2SR,12bSR)-  
N-methyl-N-(2-methylamino-1,3,4,6,7,12b-hexahydrobenzo  
[b]furo[2,3-a]quinolizin-2-yl)acetamide was reduced  
using 0.67 ml (2.28 mmol) 3.4 M sodium bis (2-methoxy-  
25 ethoxy)aluminum hydride to yield 0.082 g (48%)  
(2SR,12bSR)-2-methylamino-2-(2'-methylamino ethyl)-  
1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]  
quinolizine as a yellow oil.

30



2291S/0976A

- 48 -

17226IA

Step C: Preparation of (2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-(5',6'-dihydro-1'H-pyrimidin-2'(3'H)-one), and (2SR,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-(5',6'-dihydro-1'H-pyrimidin-2'(3'H)-one)

Into a 300 ml flask was placed 1 g (3.2 mmol) (2RS,12bSR)-2-methylamino-2-(2'-methylaminoethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine in 100 ml toluene followed by 0.95 g (5.9 mmol) 1,1'-carbonyldiimidazole. The reaction was heated to 50°C for 5 hours, after which time the reaction was treated as previously described to give 0.85 g (78%) (2SR,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-(5',6'-dihydro-1'H-pyrimidin-2'(3'H)-one) which was dissolved in ethyl acetate and ethanolic HCl was added to give the hydrochloride sesquihydrate as a white crystalline salt; m.p. 171-173°C.

Employing the above procedure but starting with the enantiomers of the diamine from Step B, there were produced the (2S,12bS)- and (2R,12bR)-enantiomers of the title compound with  $[\alpha]_D -6.2^\circ$  (C=0.0016 g/ml CH<sub>3</sub>OH) and  $[\alpha]_D +6.2^\circ$  (C=0.0016 g/ml CH<sub>3</sub>OH) respectively, m.p. 283-285°C.

In a similar manner, 0.078 g (0.25 mmol) (2SR,12bSR)-2-methylamino-2-(2'-methylaminoethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine was reacted with 0.078 gr. (0.5 mmol) 1,1'-carbonyldiimidazole to give 0.02 g (24%) (2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo

2291S/1163A

- 49 -

17226IA

[2,3-a]quinolizin)-2,4'-(5',6'-dihydro-1'H-pyrimidin-2' (3'H)-one) after heating for 24 hours at 50°C as a yellow oil.

5

EXAMPLE 28

(2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]-furo[2,3-a]quinolizin)-2,4'-(5'H-pyrimidin-2' (3'H),6' (1'H)-dione

Into a dry 50 ml 3 neck flask was placed  
10 0.164 g (0.5 mmol) (2RS,12bSR)-2-methylamino-2-(2-N-methyl acetamido)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine in 10 ml dry THF and the solution was cooled to -78°C using a dry ice/acetone bath. To this was added 0.462 ml (0.5 mmol) 1.3 M  
15 n-butyl lithium in hexane and the mixture was stirred at -78°C for 15 minutes. The solution was then transferred via cannula to a solution of 0.081 g (0.5 mmol) 1,1'-carbonyldiimidazole in 10 ml THF at 0°C and the mixture stirred 18 hours at room temperature.  
20 The solvent was removed and the residue partitioned between ethyl acetate/H<sub>2</sub>O. The layers were separated and the organic fraction washed with 3 x 50 ml H<sub>2</sub>O, 50 ml brine, dried (MgSO<sub>4</sub>) and the solvent removed. Spinning disc chromatography  
25 yielded 0.057 g (32%) (2RS,12bSR)-1',3'-dimethyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]-furo[2,3-a]quinolizin)-2,4'-(5'H-pyrimidin-2' (3'H),6' (1'H)-dione) as a yellow oil which was dissolved in ethyl acetate and ethanolic HCl added. Dilution with ether gave  
30 the hydrochloride hydrate salt as a white solid; m.p. 201-203°C.

2291S/0976A

- 50 -

17226IA

EXAMPLE 29

(2SR,12bSR)-3'-Methyl-spiro(1,3,4,6,7,12b-hexahydro-benzo[b]furo[2,3-a]quinolizine)-2,4'-(3',4',5',6'-tetrahydro-1',3'-oxazin-2'-one)

---

5                   To a 200 ml round bottomed flask with stirring bar was added 760 mg of (2RS,12bSR)-2'-methyl-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]furo [2,3-a]quinolizine)-2,3'-isoxazolidin-5'-one, 20 ml of dry THF and 9.7 ml of 1M borane-THF complex. This  
10 solution was heated at reflux under argon for 20 hours. The cooled reaction mixture was quenched with 5% aqueous HCl and partitioned between 200 ml of 5% HCl and 500 ml of ether. The layers were separated and the aqueous phase was made basic with 10N sodium  
15 hydroxide. This solution was extracted with chloroform (3 x 50 ml) and the combined chloroform extracts were washed with brine and dried over anhydrous potassium carbonate. Filtration and removal of the solvent in vacuo gave 430 mg of crude  
20 intermediate amino-alcohol. This amino-alcohol was chromatographed on 20 g of silica gel with 0.1% methanol in ammonia saturated chloroform as eluant. There was obtained 144 mg of pure intermediate amino-alcohol as a colorless foam.

25                   To a 100 ml round bottomed flask containing 100 mg of the aforementioned amino-alcohol was added 3 ml of chloroform and 406 mg of 1,1'-carbonyl-diimidazole. This mixture was stirred under argon at room temperature for 48 hours. The reaction mixture  
30 was diluted with ethyl acetate and washed with water and brine. Drying (potassium carbonate), filtration and removal of the solvent in vacuo gave an oil which was chromatographed on 5 g of silica gel with 25%

2291S/0976A

- 51 -

172261A

ethyl acetate in hexanes as eluant. The oil obtained from this chromatography was rechromatographed on 5 g of silica gel with 2% methanol in chloroform as eluant. The crystalline product thus obtained was  
5 recrystallized from a 1:1 mixture of boiling ethyl acetate-hexanes to give 69.8 mg of the title compound as white crystals, m.p. 216-218°C.

### EXAMPLE 30

10 (2RS,12bSR)-4'-Methyl-spiro(1,3,4,6,7,12b-hexahydro benzo[b]furo[2,3-a]quinolizine)-2,5'-(2',4',5',6'-  
tetrahydro-1',4'-oxazin-3'-one)

Step A: Preparation of (2RS,12bSR) 2-chloroacetoxy-  
methyl-2-(N-methyl-2-chloroacetamido)-  
15 1,3,4,6,7-12b-hexahydro-2H-benzo[b]furo  
[2,3-a]quinolizine

Chloroacetyl chloride (35 mg, 0.314 mmol) was added dropwise to a solution of (2RS, 12bSR)-2-hydroxymethyl-2-methylamino-1,3,4,6,7,12b-hexahydro-  
20 benzo[b]furo[2,3-a]quinolizine (40 mg, 0.14 mmole) and triethylamine (43  $\mu$ l, 0.314 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 ml) with stirring at 0°C, under  $\text{N}_2$ . After stirring at ambient temperature for 18 hours, the reaction mixture was treated with  $\text{H}_2\text{O}$ , the organic layer  
25 separated, washed with saturated NaCl solution, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated to dryness to give the bisacylated material, 58 mg (95%).

Step B: Preparation of (2RS,12bSR)-4'-Methyl-spiro-  
30 (1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine)-2,5'-(2',4',5',6'-tetrahydro-  
1',4'-oxazin-3'-one)

The product from Step A (58 mg, 0.13 mmol) was dissolved in toluene (15 ml) and treated with a

2291S/0976A

- 52 -

17226IA

finely ground mixture of KOH and neutral alumina (1:1) (115 mg) with stirring at ambient temperature. After 2 hours the reaction mixture was filtered, the filtrate washed with H<sub>2</sub>O, saturated NaCl solution, and dried (Na<sub>2</sub>SO<sub>4</sub>). Flash chromatography provided 33 mg (75%) of (2RS,12bSR)-4'-Methyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine)-2,5'-(2',4',5',6'-tetrahydro-1',4'-oxazin-3'-one), m.p. 159-63°C.

Anal. for C<sub>19</sub>H<sub>22</sub>N<sub>2</sub>O<sub>3</sub>.

#### EXAMPLE 31

(2RS,12bSR)-4'-Methyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine)-2,5'-piperazin-2',3'-dione

(2RS, 12bSR)-2-Aminomethyl-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine (310 mg, 1.08 mmol) and diethyl oxalate (200 µl, 1.48 mmol) in toluene (50 ml) were heated at reflux for 27 hours, then concentrated to dryness, and the residue was chromatographed to give 60 mg (16%) of (2RS,12bSR)-4'-methyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine)-2,5'-piperazin-2',3'-dione, m.p. 218-220°C.

Anal. for C<sub>19</sub>H<sub>21</sub>N<sub>3</sub>O<sub>3</sub> 1/4 H<sub>2</sub>O:  
Calc'd: C, 66.36; H, 6.30; N, 12.22.  
Found: C, 66.43; H, 6.51; N, 12.01.

#### EXAMPLE 32

Preparation of (2RS,12bSR)-4'-Methyl-spiro-(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine)-2,5'-piperazin-3'-one

2291S/0976A

- 53 -

172261A

· Step A: Preparation of (2RS, 12bSR)-2-Methylamino-1,3,4,6,7,12b-hexahydro-2-trifluoroacetamidomethyl-benzo[b]furo[2,3-a]quinolizine

Trifluoroacetic anhydride (1.84 ml, 13 mmol)  
5 dissolved in  $\text{CH}_2\text{Cl}_2$  (5 ml) was added dropwise to a solution of (2RS,12bSR)-2-aminomethyl-2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine (920 mg, 3.2 mmol) in  $\text{CH}_2\text{Cl}_2$  (50 ml) with stirring at 0°C under  $\text{N}_2$ . The reaction  
10 mixture was concentrated to dryness, and the residue was treated with  $\text{H}_2\text{O}$  and extracted with  $\text{CH}_2\text{Cl}_2$ . The aqueous acidic layer was basified with saturated  $\text{NaHCO}_3$  solution, extracted with  $\text{CH}_2\text{Cl}_2$ , the extract was washed with saturated  $\text{NaCl}$  solution and  
15 dried ( $\text{MgSO}_4$ ). Filtration and concentration to dryness provided 140 mg (12%) of the monoacylated amine.

Step B: Preparation of (2RS, 12bSR)-N-Methyl-N-(1,3,4,6,7,12b-hexahydro-2-trifluoroacetamidomethyl-benzo[b]furo[2,3-a]quinolizin-2-yl)chloroacetamide

Chloroacetyl chloride (31.8  $\mu\text{l}$ , 0.4 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 ml) was added to a solution of the  
25 product from Step A (140 mg, 0.37 mmol) and triethylamine (55.7  $\mu\text{l}$ , 0.4 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 ml) with stirring at 0°C under  $\text{N}_2$ . After stirring at ambient temperature for 2 hours, the reaction mixture was treated with ice- $\text{H}_2\text{O}$ , the organic layer  
30 was separated, washed with saturated  $\text{NaCl}$  solution and dried ( $\text{MgSO}_4$ ). Filtration and concentration to dryness gave 80 mg (47%) of the diacyl product.

2291S/0976A

- 54 -

17226IA

Step C: Preparation of (2RS,12bSR)-4'-Methyl-spiro  
(1,3,4,6,7,12b-hexahydro-2H-benzofuro[2,3-a]  
quinolizine)-2,5'-piperazin-3'-one

The product from Step B (80 mg, 0.17 mmole)  
5 and KOH (2 pellets) were stirred in methanol (20 ml)  
at ambient temperature for 2 hours. The reaction  
mixture was concentrated to dryness, then partitioned  
between CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O. The organic layer was  
separated, washed with saturated NaCl solution and  
10 dried (MgSO<sub>4</sub>). Filtration and concentration to  
dryness provided 30 mg (55%) of (2RS,12bSR)-4'-Methyl-  
spiro(1,3,4,6,7,12b-hexahydro-2H-benzofuro[2,3-a]  
quinolizine)-2,5'-piperazin-3'-one after  
chromatography. Treatment with ethanolic HCl gave  
15 the hydrochloride salt, m.p. 225-230°C.

Anal. for C<sub>19</sub>H<sub>23</sub>N<sub>3</sub>O<sub>2</sub> 2HCl 1/2 H<sub>2</sub>O:

Calcd: C, 56.03; H, 6.43; N, 10.32.

Found: C, 55.94; H, 6.43; N, 9.94.

20

EXAMPLE 33

(2SR,11bSR)-3,4,5',6,6',7,11b-octahydro-1',3'-dimethyl-  
spiro(2H-benzo[a]quinolizine)-2,4'(1'H-pyrimidin-2'-  
(3'E)-one), monohydrochloride

25

Step A: Preparation of (E,Z) 2-Carbomethoxymethyl-  
idene-1,3,4,-6,7,11b-hexahydrobenzo[a]quino-  
lizine

To a 500 ml 3-necked round bottomed flask  
with a stirring bar was added 25% potassium hydride-  
oil suspension (14.68 g, 91.52 mmol) under argon.  
30 The oil was removed by washing with hexane. The oil  
free potassium hydride was suspended in 100 ml of dry  
THF and was cooled to 0°C. Trimethylphosphonoacetate

2291S/1163A

- 55 -

172261A

(16.67 g, 91.52 mmol) in 50 ml of THF was added to the well stirred potassium hydride, dropwise over 30 minutes. When the addition was complete the mixture was allowed to stir for an additional 15 minutes then  
5 a solution of 1,2,4,6,7,11b-hexahydrobenzo[a]quinolizidin-2-one (6.14 g, 30.51 mmol) in 50 ml of THF was added dropwise, over 15 minutes. The cooling bath was allowed to expire and the mixture was stirred overnight at room temperature. The reaction mixture  
10 was diluted with ethyl acetate and washed with water (3x) and brine. Drying ( $\text{MgSO}_4$ ), filtration, and removal of the solvent in vacuo gave 7.18 g (91%) of a 1:1 mixture of (E:Z) 2-carbomethoxymethylidene-1,3,4,6,7,11b-hexahydrobenzo[a]quinolizine.

15

Step B: Preparation of (2RS,11bSR)-N-methyl-2-(2-methylamino-1,3,4,6,7,11b-hexahydrobenzo[a]quinolizidin-2-yl)acetamide

To a 125 ml pressure vessel was added a  
20 solution of (E:Z) 2-carbomethoxymethylidene-1,3,4,6,7,11b-hexahydrobenzo[a]quinolizine (7.18 g 27.90 mmol) in 25 ml of absolute ethanol. This solution was cooled to  $-78^\circ\text{C}$  and 25 ml of methylamine was condensed into the mixture. A stirring bar was  
25 added, the vessel sealed, and the mixture was stirred at room temperature for 96 hours.

The vessel was vented and the contents were concentrated in vacuo. Chromatography of the residue on 200g of silica gel using 2.5% methanol in ammonia  
30 saturated chloroform as eluant gave 5.04 g (63%) of (2RS, 11bSR)-N-methyl-2-(2-methylamino-1,3,4,6,7,11b-hexahydrobenzo[a]quinolizidin-2-yl)acetamide as a yellowish oil.



2291S/0976A

- 56 -

17226IA

Step C: Preparation of (2RS, 11bSR)-2-methylamino-2-(2-methylaminoethyl)-1,3,4,6,7,11b-hexahydrobenzo[a]quinolizine

To a 500 ml round bottomed flask with  
5 stirring bar, reflux condenser, dropping funnel and  
argon inlet was added a solution of sodium bis  
(2-methoxyethoxy)aluminum hydride (20.63 ml of a 3.4  
M solution in toluene) and 100 ml of dry THF. This  
solution was heated at reflux and a solution of the  
10 acetamide (5.04 g, 17.54 mmol) from Step B in 100 ml  
of dry THF was added dropwise over 45 minutes. When  
the addition was complete the mixture was heated at  
reflux for 2 hours. The reaction mixture was chilled  
in an ice bath and quenched with 100 ml of saturated  
15 aqueous potassium sodium tartrate solution. The  
resulting mixture was diluted with ethyl acetate (300  
ml) and the layers were separated. The organic phase  
was washed with an additional 200 ml of tartrate  
solution and 200 ml of brine. Drying ( $K_2CO_3$ ),  
20 filtration and removal of the solvent in vacuo left  
4.90 g of (2RS, 11bSR)-2-methylamino-2-(2-methylamino-  
ethyl)-1,3,4,6,7,11b-hexahydrobenzo[a]quinolizine as  
an oil.

25 Step D: Preparation of (2SR,11bSR)-1,3,4,5',6,6'7,11b-octahydro-1',3'-dimethyl-spiro(2H-benzo[a]-quinolizine)-2,4'(1'H-pyrimidin-2'(3'H)-one)-monohydrochloride

To a 200 ml round bottomed flask with  
30 stirring bar and argon inlet was added (2RS, 11bSR)-2-  
methylamino-2-(2-methylaminoethyl)-1,3,4,6,7,11b-  
hexahydrobenzo[a]quinolizine (3.57 g, 13.06 mmol), 50

2291S/0976A

- 57 -

17226IA

5 -ml of dry toluene and 1,1'-carbonyldiimidazole (3.17 g, 19.59 mmol). This solution was heated at 50°C for 20 hours. The cooled mixture was diluted with ethyl-acetate and washed with water (3 x 100 ml) and brine (200 ml). Drying (MgSO<sub>4</sub>), filtration and removal of the solvent in vacuo left an oil which was chromatographed on 150 g of silica gel using ammonia saturated chloroform as eluant. The purified free base obtained from the chromatography was  
10 crystallized as its monohydrochloride salt from ethanolic HCl. Vacuum drying at 80°C for 24 hours gave 2.36 g of the title compound, m.p. 275°C, (decomp.).

#### EXAMPLE 34

15 (2SR,12bSR)-1'-Methyl-spiro(1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin)-2,4'-(tetrahydro-(1'H)-pyrimidin-2'-one)

Step A: Preparation of (2RS,12bSR)-methyl-(2-amino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]-quinolizin-2-yl)acetate  
20

A mixture of 0.149 g (0.5 mmol) of (E,Z)-2-carbomethoxymethylidene-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine and 10 ml absolute ethanol were cooled to -78°C in a pressure vessel.  
25 Dry ammonia gas (20 ml) was condensed into the flask, which was then sealed, warmed to ambient temperature and allowed to stir for 18 hours. The pressure was released and the solvent removed in vacuo to give 0.073 g (46%) (2RS,12bSR)-methyl (2-amino-1,3,4,6,-  
30 7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)-acetate as a yellow oil after purification by spinning disc chromatography (2% methanol/ammonia saturated chloroform).

2291S/0976A

- 58 -

17226IA

Step B: Preparation of (2RS, 12bSR)-N-methyl-(2-amino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide

A mixture of 0.3 g (0.95 mmol) of (2RS, 12bSR)-methyl-(2-amino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetate and 25 ml absolute ethanol were reacted employing the procedure substantially as described in Example 27, Step A to obtain 0.25 g (84%) of (2RS, 12bSR)-N-methyl-(2-amino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide as a yellow oil.

Step C: Preparation of (2RS, 12bSR)-2-amino-2-(2-methylaminoethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine

A mixture of 0.94 ml (3.26 mmol) 3.4M sodium bis (2-methoxyethoxy)aluminum hydride and 0.25 g (0.8 mmol) of (2RS,12bSR)-N-methyl-(2-amino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)-acetamide were reacted employing the procedure substantially as described in Example 27, Step B to yield 0.146 g (61%) of (2RS,12bSR)-2-amino-2-(2'-methylaminoethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine as a yellow oil after purification by spinning disc chromatography (ammonia saturated chloroform).

Step D: Preparation of (2SR,12bSR)-1,3,4,5'6,6',7,12b-octahydro-1'-methyl-spiro(2H-benzofuro[2,3-a]quinolizine-2,4'(1'H)-pyrimidin)-2'(3'H)-one

A mixture of 0.146 g (0.49 mmol) of (2RS,-12bSR)-2-amino-2-(2'-methylaminoethyl)-1,3,4,6,7,12b-

hexahydrobenzo[b]furo[2,3-a]quinolizine and 0.158 g  
(0.98 mmol) of 1,1'-carbonyldiimidazole in 25 ml dry  
toluene were reacted employing the procedure  
substantially as described in Example 27, Step C to  
5 give 0.052 g (28%) of (2SR,12bSR)-1,3,4,5',6,6',7,-  
12b-octahydro-1'-methyl-spiro(2H-benzofuro[2,3-a]-  
quinolizine)-2,4'(1'H-pyrimidin-2'(3'H)-one)hydro-  
chloride monohydrate as a white solid; m.p. 190°C  
(dec).

10

EXAMPLE 35

(2SR,12bSR)-1,3,4,5',6,6',7,12b-octahydro-3'-methyl-  
1'-(2-hydroxyethyl)-spiro(2H-benzofuro[2,3-a]quino-  
lizine)-2,4'(1'H-pyrimidin-2'(3'H)-one)

15 Step A: Preparation of (2RS,12bSR)-methyl (2-methyl-  
amino-1,3,4,6,7,12b-hexahydrobenzo[b]furo-  
[2,3-a]quinolizin-2-yl)acetate

Employing the procedure substantially as  
described in Example 27, Step A, but stirring for only  
20 4 hours, 3 g (10 mmol) (E,Z)-2-carbomethoxymethy-  
lidine-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quino-  
lizine was reacted with 20 ml methylamine in 20 ml  
ethanol to give 2.7 g (85%) of (2RS,12bSR)-methyl-(2-  
methyldamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo-  
25 [2,3-a]quinolizin-2-yl)acetate as a yellow oil.

Step B: Preparation of (2RS,12bSR)-N-(2-propenyl)-2-  
(2-methyldamino-1,3,4,6,7,12b-hexahydrobenzo-  
[b]furo[2,3-a]quinolizin-2-yl)acetamide

30 A mixture of 2.6 g (8 mmol) of (2RS,12bSR)-  
methyl-2-(2-methyldamino-1,3,4,6,7,12b-hexahydrobenzo-  
[b]furo[2,3-a]quinolizin-2-yl)acetate, 50 ml allyl-  
amine and 50 ml absolute ethanol were refluxed for 3  
days. The reaction was cooled and the solvent removed

2291S/0976A

- 60 -

17226IA

in vacuo to give 1.8 g (63.7%) of (2RS,12bSR)-N-(2-propenyl)-2-(2-methylamino-1,2,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide as a yellow oil after medium pressure column chromatography (ammonia saturated chloroform).

Step C: Preparation of (2RS,12bSR)-2-methylamino-2-(2-(2-propenylamino)ethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine

Employing the procedure substantially as described in Example 27, Step B, 1.8 g (5 mmol) of (2RS,12bSR)-N-(2-propenyl)-2-(2-methylamino-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide and 6 ml (20 mmol) of 3.4M sodium bis (2-methoxyethoxy)aluminum hydride were reacted to give 1.2 g (70.7%) of (2RS,12bSR)-2-methylamino-2-(2-(2-propenylamino)ethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine as a yellow oil after medium pressure column chromatography (ammonia saturated chloroform).

Step D: Preparation of (2SR,12bSR)-1,3,4,5',6,6',7,12b-octahydro-1'-(2-propenyl)-3'-methylspiro(2H-benzofuro[2,3-a]quinolizin)-2,4'-(1'H-pyrimidin-2'(3'H)-one)

Employing the procedure substantially as described in Example 27, Step C, 1.2 g (3.5 mmol) of (2RS,12bSR)-2-methylamino-2-(2-(2 propenyl amino)-ethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine and 1.14 g (7 mmol) of 1,1'-carbonyl-diimidazole were reacted to give 0.8 g (63%) of (2SR,12bSR)-1,3,4,5',6,6',7,12b-octahydro-1'-(2-

2291S/0976A

- 61 -

17226IA

propenyl)-3'-methyl-spiro(2H-benzofuro[2,3-a]quino-  
lizin)-2,4'(1'H-pyrimidin-2'(3'H)-one) as a yellow  
waxy solid after purification by flask column  
chromatography, from which was made the hydrochloride  
5 dehydrate salt. m.p. 174-176°C.

Step E: Preparation of (2SR,12bSR)-1,3,4,5',6,6'-  
7,12b-octahydro-3'-methyl-1'-(2,3-dihydroxy-  
propyl)-spiro(2H-benzofuro[2,3-a]quinolizine)-  
10 2,4'(1'H-pyrimidin-2'(3'H)-one)

Employing the procedure substantially as  
described in Example 36, Step E, a mixture of 0.1 g  
(0.27 mmol) of (2SR,12bSR)-1,3,4,5',6,6',7,12b-  
octahydro-1'-(2 propenyl)-3'-methyl-spiro(2H-benzo-  
15 furo[2,3-a]quinolizin)-2,4'(1'H-pyrimidin-2'(3'H)-one),  
0.073 g (0.54 mmol) of 4-methylmorpholine-4-oxide  
monohydrate and 2 drops of osmium tetroxide (.4M  
solution in THF) were reacted to give 0.06 g (55%) of  
(2SR,12bSR)-1,3,4,5',6,6',7,12b-octahydro-3'-methyl-1'-  
20 (2,3-dihydroxypropyl)-spiro(2H-benzofuro[2,3-a]quino-  
lizin)-2,4'(1'H-pyrimidin-2'(3'H)-one) as a waxy solid  
after purification by flash column chromatography  
(ammonia saturated chloroform), from which was made  
the hydrochloride dihydrate salt. m.p. 175-177°C  
25 (dec).

Step F: Preparation of (2SR,12bSR)-1,3,4,5',6,6',-  
7,12b-octahydro-3'-methyl-1'-(2-hydroxyethyl)-  
spiro(2H-benzofuro[2,3-a]quinolizine)-2,4'-  
30 (1'H-pyrimidin-2'(3'H)-one)

A mixture of 0.125 g (0.31 mmol) of (2SR,  
12bSR)-1,3,4,5',6,6',7,12b-octahydro-3'-methyl-1'-  
(2,3-dihydroxypropyl)-spiro(2H-benzofuro[2,3-a]quino-

lizine)-2,4'(1'H-pyrimidin-2'(3'H)-one) and 0.125 ml  
20% NaOH were dissolved in 10 ml 95% ethanol and  
cooled to 0°C. To this solution was added 0.2 g  
(0.93 mmol) of sodium periodate dropwise in 5 ml  
5 H<sub>2</sub>O. The reaction was allowed to stir for 2 hours  
at 0°C, then the solvent was removed and the residue  
was partitioned between water/chloroform. The layers  
were separated, the organics dried (MgSO<sub>4</sub>) and the  
solvent removed to give the crude aldehyde which was  
10 immediately dissolved in 10 ml absolute ethanol and  
treated with a large excess ( 0.15 g) of sodium  
borohydride. After stirring 18 hours, the solvent  
was removed and the residue worked up to give 0.048 g  
(42%) of (2SR,12bSR)-1,3,4,5',6,6',7,12b-octahydro-  
15 3'-methyl-1'-(2-hydroxyethyl)-spiro(2H-benzofuro[2,3-a]  
quinolizine)-2,4'(1'H-pyrimidin-2'(3'H)-one) as a  
white crystalline solid m.p. 174-176°C.

#### EXAMPLE 36

20 (2SR,12bSR)-1,3,4,5',6,6',7,12b-octahydro-1'-methyl-  
3'-(2-hydroxyethyl)-spiro(2H-benzofuro[2,3-a]quino-  
lizine-2,4'(1'H)-pyrimidin)-2'(3'H)-one

Step A: Preparation of (2RS,12bSR)-methyl-2-(2-(2-  
propenylamino)-1,3,4,6,7,12b-hexahydro-  
25 benzo[b]furo[2,3-a]quinolizin-2-yl)acetate

A mixture of 0.148 g (0.5 mmol) of (E,Z)-  
2-carbomethoxymethylidene-1,3,4,6,7,12b-hexahydro-  
benzo[b]furo[2,3-a]quinolizin, 5 ml of allylamine and  
5 ml absolute ethanol were refluxed under a nitrogen  
30 atmosphere for 18 hours, after which the solvent was  
removed and the resulting residue purified by spinning  
disc chromatography (1:1 hexane/ammonia saturated

2291S/0976A

- 63 -

17226IA

chloroform) to give 0.063 g (36%) of (2SR,12bSR)-methyl-(2-(2-propenylamino)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl) acetate as a yellow oil.

5

Step B: Preparation of (2RS,12bSR)-N-methyl-2-(2-(2-propenyl amino)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide

A mixture of 0.86 g (2.4 mmol) of (2RS,12bSR)-methyl-2-(2-(2-propenylamino)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetate and dry methylamine were reacted employing the procedure substantially as described in Example 27, Step A to give (2RS,12bSR)-N-methyl-2-(2-(2-propenylamino)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide (0.5 g, 58%) as a yellow oil after purification by medium pressure column chromatography (ammonia saturated chloroform).

20 Step C: Preparation of (2RS,12bSR)-2-(2-propenylamino)-2-(2-methylaminoethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine

A mixture of 1.66 ml (5.6 mmol) of 3.4M sodium bis (2-methoxyethoxy)aluminum hydride and 0.5 g (1.4 mmol) of (2RS,12bSR)-N-methyl-2-(2-(2-propenylamino)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizin-2-yl)acetamide were reacted employing the procedure substantially as described in Example 27, Step B to yield 0.116 g (24%) of (2RS,12bSR)-2-(2-propenylamino)-2-(2-methylaminoethyl)-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine as a yellow oil after purification by spinning disc chromatography (3% methanol/ammonia saturated chloroform).



2291S/0976A

- 64 -

17226IA

Step D: Preparation of (2SR,12bSR)-1,3,4,5',6,6',7-  
12b-octahydro-1'-methyl-3'-(2-propenyl)-  
spiro(2H-benzofuro[2,3-a]quinolizin)-2,4'-  
(1'H-pyrimidin-2'(3'H)-one)

5 A mixture of 0.116 g (0.34 mmol) of (2RS,-  
12bSR)-2-(2-propenylamino)-2-(2-methylaminoethyl)-  
1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3-a]quinolizine  
and 0.11 g (0.68 mmol) 1,1'-carbonyldiimidazole in 20  
ml was reacted employing the procedure substantially  
10 as described in Example 27, Step C to get 0.063 g  
(51%) of (2RS,12bSR)-1,3,4,5',6,6',7,12b-octahydro-1'-  
methyl-3'-(2-propenyl)-spiro(2H-benzofuro[2,3-a]quino-  
lizin)-2,4'(1'H-pyrimidin-2'(3'H)-one) after  
purification by spinning disc chromatography (ammonia  
15 saturated chloroform), from which was made the  
hydrochloride dihydrate salt, m.p. 173-175°C (dec).

Step E: Preparation of (2SR,12bSR)-1,3,4,5',6,6',-  
7,12b-octahydro-1'-methyl-3'-(2,3-dihydroxy-  
20 propyl)-spiro-(2H-benzofuro[2,3-a]quino-  
lizine)-2,4'(1'H-pyrimidin-2'(3'H)-one)

A mixture of 0.1 g (0.27 mmol) of (2RS,-  
12bSR)-1,3,4,5',6,6',7,12b-octahydro-1'-methyl-3'-(2-  
propenyl)-spiro(2H-benzofuro[2,3-a]quinolizine)-2,4'(1'  
25 H-pyrimidin-2'(3'H)-one) and 0.73 g (.54 mmol)  
4-methylmorpholine-4-oxide monohydrate were dissolved  
in 25 ml THF, and to this was added 2 drops of a 0.4M  
solution of osmium tetroxide in THF. The reaction  
was stirred 18 hours after which time it was poured  
30 into 50 ml water and extracted with 3 x 25 ml  
chloroform. The organic layer was dried (MgSO<sub>4</sub>)  
and the solvent removed to give 0.08 g (74.4%) of  
(2SR,12bSR)-1,3,4,5',6,6',7,12b-octahydro-1'-methyl-

2291S/0976A

- 65 -

172261A

3'-(2,3-dihydroxy-propyl)-spiro(2H-benzofuro[2,3-a]quinolizine)-2,4'-(1'H-pyrimidin-2'(3'H)-one) after flash column chromatography (ammonia saturated chloroform), from which was made the hydrochloride dehydrate salt.  
5 m.p. 175-178°C.

Step F: Preparation of (2SR,12bSR)-1,3,4,5'6,6',7,-  
12b-octahydro-3'-(2-hydroxyethyl)-1'-methyl-  
spiro(2H-benzofuro[2,3-a]quinolizine)-2,4'-  
10 (1'H)-pyrimidin-2'(3'H)-one)

A mixture of 0.05 g (0.12 mmol) of (2SR,-  
12bSR)-1,3,4,5',6,6'7,12b-octahydro-1'-methyl-3'-(2,3-  
dihydroxypropyl)-spiro(2H-benzofuro[2,3-a]quinolizin)-  
2,4'-(1'H-pyrimidin-2'(3'H)-one), 0.079 g (0.36 mmol)  
15 of sodium periodate and 0.05 ml 20% NaOH are reacted  
followed by a large excess of sodium borohydride  
employing the procedure substantially as described in  
Example 35, Step F to give 0.026 g (58.6%) of (2SR,-  
12bSR)-1,3,4,5',6,6',7,12b-octahydro-3'-(2-hydroxy-  
20 ethyl)-1'-methyl-spiro(2H-benzofuro[2,3-a]quinolizine)-  
2,4'-(1'H-pyrimidin-2'(3'H)-one) as a white  
crystalline solid. m.p. 212-213°C.

#### EXAMPLE 37

25 (2SR,10bSR)-1,3,4,5'6,6',7,10b-octahydro-1',3'-  
dimethyl-spiro(2H-thieno[2,3-a]quinolizine)-2,4'-(1'H-  
pyrimidin-2'(3'H)-one)

Step A: Preparation of (E,Z)-2-carbomethoxymethyl-  
idine-1,3,4,6,7,10b-hexahydrothieno[2,3-a]-  
30 quinolizine

A mixture of 0.191 g (1 mmol) of (10bSR)-  
1,3,4,6,7,12b-hexahydrothieno[2,3-a]quinolizin-2-one,

2291S/0976A

- 66 -

172261A

0.728 g (4 mmol) of trimethylphosphonoacetate and  
0.668 g (4 mmol) 24% potassium hydride were reacted  
employing the procedure substantially as described in  
Example 24, Step A to give 0.138 g (52.3%) of (E,Z)-2-  
5 carbomethoxymethylidene-1,3,4,6,7,10b-hexahydrothieno-  
[2,3-a]quinolizine as a yellow oily solid.

Step B: Preparation of (2RS,10bSR)-N-methyl-2-(2-  
methylamino-1,3,4,6,7,10b-hexahydrothieno-  
10 [2,3-a]quinolizin-2-yl)acetamide

Employing the procedure substantially as  
described in Example 27, Step A, 0.46 g (1.7 mmol) of  
(E,Z)-2-carbomethoxymethylidene-1,3,4,6,7,10b-  
hexahydrothieno[2,3-a]quinolizine was reacted using  
15 15 ml methylamine and 10 ml ethanol to give 0.125 g  
(25%) of (2RS,10bSR)-N-methyl-2-(2-methylamino-  
1,3,4,6,7,10b-hexahydrothieno[2,3-a]quinolizin-2-yl)-  
acetamide as a brown oil after purification by medium  
pressure column chromatography.

Step C: Preparation of (2RS,10bSR)-2-methylamino-2-  
(2-methylaminoethyl)-1,3,4,6,7,10b-hexa-  
hydrothieno[2,3-a]quinolizine

A mixture of 0.125 g (0.43 mmol) of (2RS,  
25 10bSR)-N-methyl-2-(2-methylamino-1,3,4,6,7,10b-hexa-  
hydrothieno[2,3-a]quinolizin-2-yl)acetamide and 0.11  
ml (1.72 mmol) of 3.4 M sodium bis (2-methoxyethoxy  
aluminum hydride were reacted employing the procedure  
substantially described in Example 27 Step B to give  
30 0.07 g (59%) of (2RS,10bSR)-2-methylamino-2-(2-  
methylaminoethyl)-1,3,4,6,7,10b-hexahydrothieno[2,3-a]-  
quinolizine as a brown oil.

2291S/0976A

- 67 -

172261A

Step D: Preparation of (2SR,10bSR)-1,3,4,5',6,6',7,-  
10b-octahydro-1',3'-dimethyl-spiro(2H-thieno-  
[2,3-a]quinolizine)-2,4'(1'H)pyrimidin-2'-  
(3'H)-one

5           Employing the procedure substantially  
described in Example 27, Step C, 0.07 g (0.25 mmol)  
of (2RS,10bSR)-2-methylamino-2-(2-methylaminoethyl)-  
1,3,4,6,7,10b-hexahydrothieno[2,3-a]quinolizine and  
0.082 g (0.5 mmol) of 1,1'-carbonyldiimidazole were  
10 reacted to give 0.037 g (48.3%) of (2SR, 10bSR)-1,3,-  
4,5',6,6',7,10b-octahydro-1',3'-dimethyl-spiro(2H-  
thieno[2,3-a]quinolizine)-2,4'(1'H pyrimidin-2'-  
(3'H)-one) after purification by spinning disc  
chromatography, from which was made the hydrochloride  
15 monohydrate salt. m.p. 179-181°C.

#### EXAMPLE 38

(2SR,12bSR)-1,3,4,5',6,6',7,12b-octahydro-2',6'-  
dimethyl-spiro(2H-benzofuro[2,3-a]quinolizine)-2,3'-  
20 (4'H-(2H-1,2,6)thiadiazine,1',1'-dioxide)

A mixture of 0.168 g (0.54 mmol) of (2RS,-  
12bSR)-2-methylamino-2-(2-methylaminoethyl)-1,3,4,6,7,-  
12b-hexahydrobenzo[b]furo[2,3-a]quinolizine and 0.109  
g (1.08 mmol) of triethylamine in 10 ml methylene  
25 chloride was cooled to -78°C under N<sub>2</sub>. Sulfuryl  
chloride (0.144 g, 1.08 mmol) was reacted employing  
the procedure substantially as described in Example  
12 to give 0.062 g (28%) of (2SR,12bSR)-1,3,4,5',6,-  
6',7,12b-octahydro-2',6'-dimethyl-spiro(2H-benzofuro-  
30 [2,3-a]quinolizine)-2,3'-(4'H-(2H-1,2,6)thiadiazine,-  
1',1'-dioxide)hydrochloride m.p. 250°C (dec).

2291S/0976A

- 68 -

17226IA

EXAMPLE 39

(2SR,12bSR)-1,3,4,6,7,12b-hexahydro-spiro-(2H-benzothieno[2,3-a]quinolizin)-2,5'-(oxazolidin-2'-one)

5      Step A: Preparation of (2SR,12bSR)-2-aminomethyl-2-hydroxy-1,3,4,6,7,12b-hexahydrobenzo[b]-thieno[2,3-a]quinolizine

Into a 500 ml flask was placed 0.8 g (3.11 mmol) of 1,3,4,6,7,12b-hexahydrobenzo[b]thieno[2,3-a]quinolizin-2-one in 200 ml wet THF which had  
10      previously been saturated with methylamine at 0°C. To this mixture was added 1 g (6.21 mmol) of diethyl cyanophosphonate. The reaction was then carried out employing the procedure substantially as described in Example 16, Step A to give (2SR,12bSR)-2-aminomethyl-  
15      2-hydroxy-1,3,4,6,7,12b-hexahydrobenzo[b]thieno[2,3-a]quinolizine which was carried on without further purification.

20      Step B: Preparation of (2SR,12bSR) (1,3,4,6,7,12b-hexahydro-spiro-(benzo[b]thieno[2,3-a]quinolizin)-2,5'-(oxazolidin-2'-one)

A mixture of amino alcohol from Step A and 100 ml of toluene was treated with 0.645 g (3.98 mmol) 1,1'-carbonylidiimidazole employing the procedure  
25      substantially as described in Example 1, Step G to obtain 0.2 g (21%) (2SR,12bSR) (1,3,4,6,7,12b-hexahydro-spiro-(benzo[b]thieno[2,3-a]quinolizin)-2,5'-(oxazolidin-2'-one) from which was made the hydrochloride salt. m.p. 240°C (dec).

30

EXAMPLE 40

(2SR,12bSR)-3'-Methyl-spiro(1,3,4,6,7,12b-hexahydro-benzo[b]thieno[2,3-a]quinolizin)-2,5'-(oxazolidin-2'-one)

- 5           Employing the procedure described in Example 3, (2SR,12SR)-spiro (1,3,4,6,7,12b-hexahydrobenzo[b]-thieno[2,3-a]quinolizin)-2,5' (oxazolidin-2'-one) (Example 39) (35 mg, 0.1 mmole) gave 21 mg (57%) of the title compound as the hydrochloride, m.p. 280°C.

10

EXAMPLE 41

(2RS,12bSR)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo[2,3-a]quinolizine)-2,5'-(pyrrolidin-2'-one)

- 15   Step A: Preparation of (2RS,12bSR)-Ethyl-2-methyl-acetamido-1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo[2,3-a]quinolizine-2-carboxylate  
          (2RS,12bSR)-Ethyl-2-methylamino-1,3,4,6,7,12,-  
20   b-hexahydro-2H-benzo[b]furo[2,3-a]quinolizine-2-carboxylate (Step B, Example 19) was dissolved in 5 ml of methylene chloride and 50 mg (0.49 mmol, 0.07 ml) of triethylamine was added followed by 39 mg (0.5 mmol, 0.035 ml) of acetyl chloride. After 6 hours at  
25   room temperature, this was diluted with saturated sodium bicarbonate and the product filtered and concentrated. The residue obtained was chromatographed (SiO<sub>2</sub>,NH<sub>3</sub> saturated chloroform) to give 82 mg of the title compound as an oil. M+=370.

30

2291S/0976A

- 70 -

17226IA

Step B: Preparation of (2RS,12bSR)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo[2,3-a]quinolizine)-2,2'-(pyrrolidin-3',5'-dione)

5 (2RS,12bSR)-Ethyl-2-methylacetamido-1,3,4,-  
6,7,12b-hexahydro-2H-benzo[b]furo[2,3-a]quinolizine-  
2-carboxylate 36 mg (0.102 mmol) was dissolved in 2  
ml of DMSO and 23 mg (0.203 mmol) of potassium-  
t-butoxide was added. After 1.5 hour at room  
10 temperature the reaction was quenched with saturated  
NH<sub>4</sub>Cl solution, made basic and washed with ethyl  
acetate. The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>),  
filtered and concentrated to give an oil which was  
chromatographed (SiO<sub>2</sub>, 10% MeOH/CHCl<sub>3</sub>) to give  
15 the title compound. M+(324).

Step C: Preparation of (2RS,12bSR,3'RS)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo[2,3-a]quinolizine)-2,2'-(3'-hydroxy pyrrolidin-5'-one)

20 (2RS,12bSR)-1'-Methyl-spiro-(1,3,4,6,7,12b-  
hexahydro-2H-benzo[b]furo[2,3-a]quinolizine)-2,2'-  
(pyrrolidin-3',5'-dione) 32 mg (0.1 mmol) was  
dissolved in 5 ml of ethanol and cooled to 0°C. To  
25 this was added 4 mg of NaBH<sub>4</sub>. After 1 hour this  
was diluted with water and extracted with ethyl  
acetate which was dried (Na<sub>2</sub>SO<sub>4</sub>) filtered and  
concentrated to give 27 mg of the subject compound as  
a mixture of alcohols. M+(326).

30

2291S/0976A

- 71 -

172261A

Step D: Preparation of (2RS,12bSR,3'RS)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]-furo[2,3-a]quinolizine)-2,2'-(3'-mesyloxy pyrrolidin-5'-one)

5 To 25 mg (0.076 mmol) of (2RS,12bSR,3'RS)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo-[2,3-a]quinolizine)-2,2'-(3'-hydroxy pyrrolidin-5'-one) dissolved in 5 ml of methylene chloride was added 10 mg (0.099 mmol, 0.014 ml) of triethylamine  
10 followed by 11 mg (0.099 mmol, 0.008 ml) of methanesulfonyl chloride. After 1 hour at 0°C the solution was diluted with saturated NaHCO<sub>3</sub> solution and washed with methylene chloride which was dried (Na<sub>2</sub>SO<sub>4</sub>) filtered and concentrated to give 27 mg  
15 (89%) of the title product which as a mixture of mesylates.

Step E: Preparation of (2RS,12bSR,)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo-[2,3-a]quinolizine)-2,2'-(pyrrolid-3',4'-en-5'-one)

20 To a solution of (2RS,12bSR,3'RS)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo[2,3-a]quinolizine)-2,2'-(3'-meslyloxy pyrrolidin-5'-one)  
25 (32 mg, 0.076 mmol) in 2 ml of methylene chloride was added 12 mg (0.026 mmol) of 1,8-diazabicyclo[5.4.0]-undec-7-ene and the solution was warmed to 40°C. After 30 minutes the reaction was diluted with methylene chloride, washed with water, dried  
30 (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to a residue which was used directly in Step F. M+(308).



2291S/1163A

- 72 -

17226IA

Step F: Preparation of (2RS,12bSR)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo-[2,3-a]quinolizine)-2,5'-(pyrrolidin-2'-one)

- To a solution of (2RS,12bSR)-1'-Methyl-spiro-(1,3,4,6,7,12b-hexahydro-2H-benzo[b]furo[2,3-a]-quinolizine)-2,2'-(pyrrolid-3',4'-en-5'-one) (59 mg, 0.19 mmol) in 5 ml of ethanol was added 25 mg of 5% Palladium on charcoal and the reaction was charged with an atmosphere of hydrogen at 1 atmosphere.
- After 1.5 hour the reaction was filtered through celite, the filtrate was evaporated and the residue was chromatographed ( $\text{SiO}_2$ , 10% MeOH/ $\text{CHCl}_3$ ) to yield 39 mg of the title compound. m.p. 189-193°C. HCl salt trihydrate (MeOH) M+(310).

15

#### EXAMPLE 42

1,1',2'3,3'4,4',6,7,12b-Decahydro-4'-methanesulfonyl-spiro(2H-benzofuro(2,3-a)quinolizine-2,5'(5H-1,4-diazepin-7'(6'H)-one) monohydrochloride

- Step A: Preparation of 1,1',2',3,3',4,4',6,7,12b-decahydro-spiro-(2H-benzofuro(2,3-a)quinolizine)-2,5'-(5H-1,4-diazepin-7'(6'H)-one)
- To a 10 ml round bottomed flask with stirring bar and argon inlet was added (E,Z)-2-carbomethoxy-methylidene-1,3,4,6,7,12b-hexahydrobenzo[b]furo[2,3,-a]quinolizine (900 mg, 3.02 mmol), 4.5 ml of methanol and ethylenediamine (8.09 g, 134.6 mmol), freshly distilled from calcium hydride. This mixture was stirred at room temperature for 40 hours, then diluted with ethylacetate and washed with water and brine. Drying ( $\text{Na}_2\text{SO}_4$ ), filtration, removal of the solvent in vacuo followed by chromatographed of the residue on 25 g of silica gel using 6% methanol

2291S/0976A

- 73 -

17226IA

in chloroform as eluant provided 550 mg of intermediate diazepinone.

5      Step B: Preparation of 1,1',2'3,3'4,4',6,7,12b-decahydro-4'-methanesulfonyl-spiro-(2H-benzofuro(2,3-a)quinolizine)-2,5'-(5H-1,4-diazepin-7'(6'H)-one) monohydrochloride

To a 25 ml round bottomed flask containing intermediate diazepinone from Step A (396 mg, 1.21 mmol) was added 30 ml of dry methylene chloride and diisopropylethylamine (2.12 ml, 12.17 mmol). The resulting solution was cooled to 0°C in an ice bath and methanesulfonyl chloride (0.47 ml, 6.07 mmol) was added in one portion. The mixture was stirred at 0°C for 30 minutes then diluted with methylene chloride and washed with water and brine. Drying (Na<sub>2</sub>SO<sub>4</sub>), filtration, removal of the solvent in vacuo, followed by chromatography of the residue on 100 g of silica gel using 5% methanol in chloroform as eluant provided 134 mg of the title compound free base. The hydrochloride salt was crystallized from ethanolic HCl and dried in vacuo. m.p. 282-285°C.

#### EXAMPLE 43

#### 25      Pharmaceutical Formulation

<u>Ingredient</u>	<u>Mg/Capsule</u>
(2RS,12bSR)-3'-methyl-spiro	
(1,3,4,6,7,12b-hexahydro-	
30 benzo[b]furo[2,3-a]quinolizine)-	
2,4'-imidazolidin-2'-one	6
starch	87
magnesium stearate	7

2291S/0976A

- 74 -

17226IA

- The active ingredient, starch and magnesium stearate are blended together. The mixture is used to fill hard shell capsules of a suitable size at a fill weight of 100 mg per capsule.

5

EXAMPLE 44

	<u>Ingredient</u>	<u>Mg/Capsule</u>
10	(2SR,12bSR)-1',3'-dimethyl-spiro (1,3,4,6,7,12b-hexahydro-benzo[b]- furo[2,3-a]quinolizine)-2,4'- (5',6'-dihydro-1',3'-diazin-2'-one)	6
	starch	87
	magnesium stearate	7

15

20


25

30

WHAT IS CLAIMED IS:

1. A compound of structural formula I

5



10

wherein

Ar represents - R<sup>1</sup>, R<sup>2</sup>-benzo or a heterocyclic ring structure of aromatic character which is selected from R<sup>1</sup>, R<sup>2</sup>-benzo[b]furo, R<sup>1</sup>, R<sup>2</sup>-benzo[b]thieno, thieno, furo, R<sup>1</sup>, R<sup>2</sup>-pyridino, thiazolo, imidazo, and pyrazolo, wherein

R<sup>1</sup> and R<sup>2</sup> are independently:

- 1) hydrogen,
- 2) halo,
- 3) hydroxy,
- 4) C<sub>1-3</sub>alkoxy, or
- 5) C<sub>1-6</sub>alkyl;

20

B represents a spiroheterocycle of 4-7 members with up to 2 heteroatoms one of which is bonded to the spiro carbon, and if Ar is benzo, that heteroatom is nitrogen, wherein the members are independently

25


-CH<sub>2</sub>-, -CH-, -C-, -C-, -C-, -O-, -NR<sup>3</sup>-,  
 $\begin{array}{c} | \\ R^3 \end{array}$   $\begin{array}{c} || \\ O \end{array}$   $\begin{array}{c} || \\ S \end{array}$   $\begin{array}{c} || \\ N-CN \end{array}$

30

or -SO<sub>2</sub>- wherein


R<sup>3</sup> is 1) hydrogen,


h

- 2)  $\overset{\text{O}}{\parallel}\text{-C-R}$ , wherein R is hydrogen or  $\text{C}_{1-3}$  alkyl,
- 3)  $\text{C}_{1-6}$  alkyl, either unsubstituted or substituted with one or more of;
- hydroxy,
  - carboxy,
  - $\text{C}_{1-3}$ alkoxycarbonyl,
  - halo,
  - $\text{C}_{1-3}$ alkoxy,
  - $\text{-CONR}^6\text{R}^7$ , wherein  $\text{R}^6$  and  $\text{R}^7$  have independently from each other the meaning of hydrogen and  $\text{C}_{1-5}$ alkyl, or  $\text{R}^6$  and  $\text{R}^7$  form together with the nitrogen atom, to which they are bonded, a 5-7 membered ring or they form together with the nitrogen atom to which they are bonded a 6-membered ring which comprises a further heteroatom selected from O, N and S,
  - $\text{-NR}^6\text{R}^7$ , wherein  $\text{R}^6$  and  $\text{R}^7$  are as defined above,
  - 
 $\text{C}_6\text{H}_4(\text{R}^1)(\text{R}^2)$ , wherein  $\text{R}^1$  and  $\text{R}^2$  is as defined above.
  - $\text{-SO}_2\text{NR}^6\text{R}^7$ , wherein  $\text{R}^6$  and  $\text{R}^7$  is as defined above or
  - $\text{-SO}_2(\text{C}_{1-3}\text{alkyl})$ ; and
- $\text{R}^5$  is 1) hydrogen,

- 2)  $C_{1-6}$  alkyl, either unsubstituted or substituted with one or more groups having the formulae  $-OR^8$ ,  $-NR^8COR^8$ , or  $-CO_2R^8$ ,  
 5 in which formulae  $R^8$  is hydrogen or alkyl having 1-6 carbon atoms,  
 3)  $-CO_2R^8$ , wherein  $R^8$  is hydrogen or alkyl having 1-6 carbon atoms, or  
 4)  $-CONR^6R^7$ , wherein  $R^6$  and  $R^7$  are as defined above, and  
 10 salts of the compounds of formula I.

2. Compounds according to claim 1, wherein the spiro-heterocycle is selected from groups having the  
 15 structure





wherein  $R^3$  is as defined in claim 1  
and salts of said compounds of formula I.

3. Compounds according to claim 1 or 2,

wherein  
Ar is  $R^1$ ,  $R^2$ -benzo[b]furo,  $R^1R^2$ -benzo[b]thieno, thieno,  
benzo or furo  
or salts of said compounds of formula I.

4. Compounds according to one of the claims

1 - 3 wherein  
Ar is  $R^1$ ,  $R^2$ -benzo[b]furo,  $R^1$ ,  $R^2$ -benzo[b]thieno- or  
benzo-; wherein  
 $R^1$  and  $R^2$  are hydrogen or halo,  
B is a spiro-imidazolidin-2-one or spiro-(5,6-dihydro-  
1H-pyrimidin-2(3H)-one: wherein  
 $R^3$  is an alkyl group having 1-6 carbon atoms and  
 $R^5$  is hydrogen or alkyl having 1-6 carbon atoms or  
salts of the compounds of formula I.

5. Compounds according to claim 4 characterized in that

Ar is R<sup>1</sup>, R<sup>2</sup>-benzo[b]furo- wherein

R<sup>1</sup> and R<sup>2</sup> are hydrogen,

5 R<sup>3</sup> is methyl and

R<sup>5</sup> is hydrogen

or salts of the compounds of formula I.

6. Compounds according to claim 5 characterized in that they are the

(2R, 12bS)-1',3'-dimethylspiro(1,3,4,6,7,12b-hexahydro-benzo[b]furo-[2,3-a]quinolizin)-2,4'-imidazolidin-2'-one, or salts thereof or the

15 (2S, 12bS)-1',3'-dimethylspiro-(1,3,4,6,7,12b-hexahydro-benzo[b]furo[2,3-a]quinolizin)-2,4'-(5',6'-dihydro-1'H-pyrimidine-2'(3'H)-one), or salts thereof.

7. Compounds according to one of the claims 1 - 6 characterized in that they have a configuration in  
20 which the heteroatom in ring B attached to carbon 2, and the hydrogen at 12b are trans as well as the enantiomers of said compounds and furthermore salts of said compounds.

25 8. Pharmaceutical composition for antagonizing the  $\alpha_2$ -adrenoceptor characterized in that they contain as  $\alpha_2$ -adrenoceptor antagonist at least one compound of formula I according to claim 1 or a pharmaceutically acceptable salt of said compound.


30

9. Pharmaceutical composition according to claim 8 characterized in that it contains as pharma-




ceutically effective ingredient at least one compound according to one of the claims 2 - 7 or a pharmaceutically effective salt of said compounds.


- 5 10. Process for the preparation of compounds of formula I




wherein the substituents have the same meaning as defined in claim 1  
 15 and salts of the compounds of formula I characterized in that corresponding compounds of formula I in which the spiroheterocycle B have the structure




are prepared by treating a starting material of formula II  
 30




5 in which the partial structure of formula



10 is the partial structure




15 or the partial structure




or the partial structure

20




25 or the partial structure




30

or the partial structure




5 (14a)


with carbonyldiimidazole or with phosgene and that optionally corresponding products of formula I in which  $R^3$  is hydrogen are further alkylated, benzylated or acylated to yield corresponding compounds of formula I wherein  $R^3$  is the acyl group or the optionally substituted alkyl group stated in claim 1, or that a compound of formula I is prepared, in which the spiroheterocycle B has the structure



or the structure




25 or the structure




by treating a corresponding starting material of


formula II in which the partial structure of formula



has the formula




with thiocarbonyldiimidazole or thiophosgene to yield the compounds of formula I in which the spiroheterocycle has the structure (2) or with SO<sub>2</sub>Cl<sub>2</sub> to yield the compounds of formula I in which the spiroheterocycle has the structure (3) or with (COC<sub>2</sub>H<sub>5</sub>)<sub>2</sub> to yield the compounds in which the spiroheterocycle has the structure (16) and that optionally resulting compounds of formula I in which the spiroheterocycle has the structure (2) or (3) or (16) wherein at least one of the radicals R<sup>3</sup> has the meaning of a hydrogen atom are further acylated or alkylated respectively benzylated to yield corresponding compounds of formula I wherein at least one of the radicals R<sup>3</sup> is the acyl group or the optionally substituted alkyl group defined in claim 1 or the compounds of formula I are prepared, in which the spiroheterocycle of formula B is a group having the structure



Dr. IM.-vd  
20.5.1986


by reacting a starting material of formula II, in which the partial structure of formula




has the following structure




with  $[(CH_3)_2N]_2P(=O)OCH=CH_2$  and  $M-C_4H_9Li$  or that compounds of formula I are prepared in which the spiro-heterocycle has the structure




20 or the structure




25 by reacting a starting material of formula II in which the partial structure



has the formula




5 or the formula



10


with  $\text{R}^3\text{-NCO}$  and that the resulting compounds of formula I in which one of the radicals  $\text{R}_3$  is hydrogen are optionally further acylated or alkylated respectively benzylated to yield corresponding compounds of formula I in which the radical  $\text{R}^3$  is the acyl group or the optionally substituted alkyl group defined in claim 1, or the compounds of formula I are prepared in which the spiroheterocycle B has the structure

20




or the structure

25




30 or the structure



Dr.IM.-vd  
20.5.1986

by treating a starting material of formula II in which the partial structure of formula




has the formula




either with  $\text{R}^3\text{NH NHR}^3$ , or with  $\text{R}^3\text{NH}(\text{CH}_2)_2\text{NHR}^3$   
or with  $\text{HONHR}^3$


and that optionally resulting compounds of formula I  
15 in which  $\text{R}^3$  is hydrogen, are further acylated or  
alkylated respectively benzylated to yield the compounds  
of formula I wherein  $\text{R}^3$  is the acyl group or the optio-  
nally substituted alkyl group defined in claim 1,  
or that a compound of formula I is prepared in which  
20 the spiroheterocycle B has the structure




25 by treating a starting material of formula II  
wherein the partial structure of formula



has the structure




- 5 with KOH and neutral alumina and wherein optionally  
 resulting compounds of formula I wherein  $R^3$  is hydrogen  
 are further acylated or alkylated respectively benzylated  
 to yield the corresponding compounds of formula I where-  
 in  $R^3$  is the acyl radical or the optionally substituted  
 10 alkyl radical defined in claim 1,  
 or that a compound of formula I is prepared in which  
 the spiroheterocycle B has the structure



(17)

by treating a starting material of formula II in which  
 the partial structure



has the formula




- 25 with KOH and that optionally corresponding compounds of  
 30 formula I in which one of the radicals  $R^3$  is a hydrogen  
 are further acylated or alkylated respectively benzylated  
 to produce corresponding compounds of formula I wherein




the radical  $R^3$  is an acyl group or an optionally substituted alkyl group as defined in claim 1,  
 or that compounds of formula I are prepared in which  
 the spiroheterocycle B has the structure

5




10 by preparing first a compound of formula I in which  
 the spiroheterocycle B has the structure

15




and treating said compound with hydrogen in the presence  
 of a noble metal catalyst,  
 and that optionally the resulting compounds in which  
 20  $R^3$  is hydrogen are further acylated or alkylated re-  
 spectively benzylated to yield compounds of formula I  
 in which  $R^3$  is the acyl group or the optionally substituted  
 alkyl group defined in claim 1,  
 or that compounds of formula I wherein the spirohetero-  
 25 cycle B has the formula



30 are prepared by cyclizing the starting material of  
 formula II wherein the partial structure of formula



5 is a group of formula



10 or a derivative of said carboxylic acid, and that  
the resulting compounds of formula I wherein  $R^3$  is  
hydrogen, are optionally further acylated or alkylated  
respectively benzylated yielding corresponding com-  
pounds of formula I in which  $R^3$  is an acyl group or  
15 an optionally substituted alkyl group as defined in  
claim 1,  
and that all the prepared compounds of formula I are  
isolated in the free form or as salts thereof.

20 11. Process according to claim 10,  
characterized in that a compound of formula I accord-  
ing to one of the claims 2 - 7 or a salt thereof is  
prepared.

25

30

Dr.IM.-vd  
20.5.1986

M

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 204 254**  
**A3**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 86107199.1

(22) Date of filing: 27.05.86

(51) Int. Cl.: **C 07 D 491/22, C 07 D 495/22,**  
**C 07 D 513/22, C 07 D 498/22,**  
**C 07 D 471/20, C 07 D 471/22,**  
**A 61 K 31/435, A 61 K 31/505,**  
**A 61 K 31/50, A 61 K 31/535,**  
**A 61 K 31/55 // (C07D491/22,**  
**307:00, 235:00, 221:00, 221:00),**  
**(C07D491/22, 307:00, 307:00,**  
**221:00, 221:00), (C07D491/22,**  
**307:00, 231:00, 221:00, 221:00)**

(30) Priority: 03.06.85 US 740609

(43) Date of publication of application: 10.12.86  
Bulletin 86/50

(84) Designated Contracting States: AT BE CH DE FR GB IT LI  
LU NL SE

(88) Date of deferred publication of search  
report: 02.03.88 Bulletin 88/9


(71) Applicant: MERCK & CO. INC., 126, East Lincoln Avenue  
P.O. Box 2000, Rahway New Jersey 07065-0900 (US)

(72) Inventor: Baldwin, John J., 621 Gypsy Hill Circle,  
Gwynedd Valley, PA 19437 (US)  
Inventor: Huff, Joel R., 738 Bergey Mill Road, Ledersach,  
PA 19450 (US)  
Inventor: Vacca, Joseph P., 766 Eisenhower Drive,  
Telford, PA 18969 (US)  
Inventor: Young, Steven D., J-10 Forge Gate Apts.  
Snyder Road, Lansdale, PA 19446 (US)  
Inventor: De Solms, Jane, 735 Port Indian Road,  
Norristown, PA 19403 (US)  
Inventor: Guare, James P., Jr., R.D. 2 Kumry Road,  
Quakertown, PA 18951 (US)

(74) Representative: Blum, Rudolf Emil Ernst et al, c/o E.  
Blum & Co Patentanwälte Vorderberg 11,  
CH-8044 Zürich (CH)

(54) Substituted hexahydro arylquinolizines, processes for their preparation and pharmaceutical compositions containing them.

(57) Compounds of formula



wherein

Ar represents R<sup>1</sup>, R<sup>2</sup>-benzo or a heterocyclic ring structure of aromatic character which is selected from R<sup>1</sup>, R<sup>2</sup>-benzo[b]furo, R<sup>1</sup>, R<sup>2</sup>-benzo[b]thieno, thieno, furo, R<sup>1</sup>, R<sup>2</sup>-pyridino, thiazolo, imidazo, and pyrazolo, and B represents a spiroheterocycle of 4-7 members with up to 2 heteroatoms one of which is bonded to the spiro carbon, are selective  $\alpha_2$ -adrenergic receptor antagonists and thereby useful as antidepressants, antihypertensives, ocular antihypertensives, antidiabetics, platelet aggregation inhibitors, antiobesity agents, and modifiers of gastrointestinal motility.

**EP 0 204 254 A3**



European Patent  
Office

# EUROPEAN SEARCH REPORT

0204254

Application Number

EP 86 10 7199

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.4)
A	FR-A-1 524 953 (KONINKLIJKE PHARMACEUTISCHE FABRIEKEN VAN HET BROCADES-STHEEMAN & PHARMACIA) -----		C 07 D 491/22 C 07 D 495/22 C 07 D 513/22 C 07 D 498/22 C 07 D 471/20 C 07 D 471/22 A 61 K 31/435 A 61 K 31/505 A 61 K 31/50 A 61 K 31/535 A 61 K 31/55 // (C 07 D 491/22 C 07 D 307:00 C 07 D 235:00 C 07 D 221:00 C 07 D 221:00 ) (C 07 D 491/22 C 07 D 307:00
			TECHNICAL FIELDS SEARCHED (Int. CL.4)
			-/-
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 13-11-1987	Examiner VAN GEYT J.J.A.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document			



European Patent  
Office

# EUROPEAN SEARCH REPORT

Page 0204254  
Application Number

EP 86 10 7199

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
			C 07 D 307:00 C 07 D 221:00 C 07 D 221:00 ) (C 07 D 491/22 C 07 D 307:00 C 07 D 231:00 C 07 D 221:00 C 07 D 221:00 ) (C 07 D 491/22 C 07 D 307:00 C 07 D 221:00 ) C 07 D 221:00
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 13-11-1987	Examiner VAN GEYT J.J.A.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

RPO FORM 1503 03.82 (P0401)

12

# EUROPEAN PATENT APPLICATION

21 Application number: 86107199.1

22 Date of filing: 27.05.86

51 Int. Cl.: **C 07 D 491/22**, C 07 D 495/22,  
 C 07 D 513/22, C 07 D 498/22,  
 C 07 D 471/20, C 07 D 471/22,  
 A 61 K 31/435, A 61 K 31/505,  
 A 61 K 31/50, A 61 K 31/535,  
 A 61 K 31/55 // (C07D491/22,  
 307:00, 235:00, 221:00, 221:00),  
 (C07D491/22, 307:00, 307:00,  
 221:00, 221:00), (C07D491/22,  
 307:00, 231:00, 221:00, 221:00)

30 Priority: 03.06.85 US 740609

43 Date of publication of application: 10.12.86  
 Bulletin 86/50

64 Designated Contracting States: AT BE CH DE FR GB IT LI  
 LU NL SE

68 Date of deferred publication of search  
 report: 02.03.88 Bulletin 88/9


71 Applicant: MERCK & CO. INC., 126, East Lincoln Avenue  
 P.O. Box 2000, Rahway New Jersey 07065-0900 (US)

72 Inventor: Baldwin, John J., 621 Gypsy Hill Circle,  
 Gwynedd Valley, PA 19437 (US)  
 Inventor: Huff, Joel R., 738 Bergey Mill Road, Ledereach,  
 PA 19450 (US)  
 Inventor: Vacca, Joseph P., 766 Eisenhower Drive,  
 Telford, PA 18969 (US)  
 Inventor: Young, Steven D., J-10 Forge Gate Apts.  
 Snyder Road, Lansdale, PA 19446 (US)  
 Inventor: De Solms, Jane, 735 Port Indian Road,  
 Norristown, PA 19403 (US)  
 Inventor: Guare, James P., Jr., R.D. 2 Kumry Road,  
 Quakertown, PA 18951 (US)

74 Representative: Blum, Rudolf Emil Ernst et al, c/o E.  
 Blum & Co Patentanwälte Vorderberg 11,  
 CH-8044 Zürich (CH)

54 Substituted hexahydro arylquinolizines, processes for their preparation and pharmaceutical compositions containing them.

57 Compounds of formula



wherein

Ar represents R<sup>1</sup>, R<sup>2</sup>-benzo or a heterocyclic ring structure of aromatic character which is selected from R<sup>1</sup>, R<sup>2</sup>-benzo[b]furo, R<sup>1</sup>, R<sup>2</sup>-benzo[b]thieno, thieno, furo, R<sup>1</sup>, R<sup>2</sup>-pyridino, thiazolo, imidazo, and pyrazolo, and B represents a spiroheterocycle of 4-7 members with up to 2 heteroatoms one of which is bonded to the spiro carbon, are selective α<sub>2</sub>-adrenergic receptor antagonists and thereby useful as antidepressants, antihypertensives, ocular antihypertensives, antidiabetics, platelet aggregation inhibitors, antiobesity agents, and modifiers of gastrointestinal motility.

EP 0 204 254 A3



European Patent  
Office

# EUROPEAN SEARCH REPORT

0204254

Application Number

EP 86 10 7199

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	FR-A-1 524 953 (KONINKLIJKE PHARMACEUTISCHE FABRIEKEN VAN HET BROCADES-STHEEMAN & PHARMACIA) -----		C 07 D 491/22 C 07 D 495/22 C 07 D 513/22 C 07 D 498/22 C 07 D 471/20 C 07 D 471/22 A 61 K 31/435 A 61 K 31/505 A 61 K 31/50 A 61 K 31/535 A 61 K 31/55 // (C 07 D 491/22 C 07 D 307:00 C 07 D 235:00 C 07 D 221:00 C 07 D 221:00 ) (C 07 D 491/22 C 07 D 307:00
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			-/-
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 13-11-1987	Examiner VAN GEYT J. J. A.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	



European Patent  
Office

# EUROPEAN SEARCH REPORT

Page 0204254

Application Number

EP 86 10 7199

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
			C 07 D 307:00 C 07 D 221:00 C 07 D 221:00 ) (C 07 D 491/22 C 07 D 307:00 C 07 D 231:00 C 07 D 221:00 C 07 D 221:00 ) (C 07 D 491/22 C 07 D 307:00 C 07 D 221:00 ) C 07 D 221:00
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 13-11-1987	Examiner VAN GEYT J.J.A.
<b>CATEGORY OF CITED DOCUMENTS</b>			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03/82 (1/04/01)



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**